arXiv:0809.0808v1 [math.FA] 4 Sep 2008

Characteristic Classes on Grassmann Manifolds

Jianwei Zhou and Jin Shi

Department of Mathematics, Suzhou University, Suzhou 215006, P. R. China
Suzhou Senior Technical Institute, Suzhou 215007, P.R. China

Abstract

In this paper, we use characteristic classes of the canonical vector bundles and the Poincaré dualality to study the structure of the real homology and cohomology groups of oriented Grassmann manifold $G(k,n)$. Show that for $k=2$ or $n\leq 8$, the cohomology groups $H^*(G(k,n),\mathbf{R})$ are generated by the first Pontrjagin class, the Euler classes of the canonical vector bundles. In these cases, the Poincaré dualality: $H^q(G(k,n),\mathbf{R})\to H_{k(n-k)-q}(G(k,n),\mathbf{R})$ can be given explicitly.

§1. Introduction

Let $G(k,n)$ be the Grassmann manifold formed by all oriented k-dimensional subspaces of Euclidean space $\mathbf{R}^n$. For any $\pi\in G(k,n)$, there are orthonormal vectors $e_1,\cdots,e_k$ such that π can be represented by $e_1\wedge\cdots\wedge e_k$. Thus $G(k,n)$ becomes a submanifold of the space $\bigwedge^k(\mathbf{R}^n)$, thus we can use moving frame to study the Grassmann manifolds.

There are two canonical vector bundles $E=E(k,n)$ and $F=F(k,n)$ over $G(k,n)$ with fibres generated by vectors of the subspaces and the vectors orthogonal to the subspaces respectively. Then we have Pontrjagin classes $p_i(E)$ and $p_j(F)$ with relationship

$$(1+p_1(E)+\cdots)(1+p_1(F)+\cdots)=1.$$

If k or $n-k$ is even number, we have Euler classes $e(E)$ or $e(F)$.

Key words and phrases. Grassmann manifold, fibre bundle, characteristic class, homology group, Poincaré duality.

Subject classification. 14M15, 55R10, 55U30, 57T15.

E-mail: jwzhou@suda.edu.cn and ttwomei@126.com

The oriented Grassmann manifolds are classifying spaces for oriented vector bundles. For any oriented vector bundle τ: $\xi \to M$ with fibre type $\mathbf{R}^k$, there is a map g: $M \to G(k,n)$ such that ξ is isomorphic to the induced bundle $g^*E(k,n)$. If the maps g_1, g_2: $M \to G(k,n)$ are homotopy, the induced bundles $g_1^*E(k,n)$ and $g_2^*E(k,n)$ are isomorphic. Then the characteristic classes of the vector bundle τ are the pullback of the characteristic classes of the vector bundle $E(k,n)$.

In this paper, we use characteristic classes and the Poincaré dualality to study the real homology and cohomology groups of oriented Grassmann manifolds. The characteristic classes of the canonical vector bundles can be represented by curvatures and are harmonic forms on the Grassmann manifolds, see [7], [8], [13]. For $k=2$ or $n \leq 8$, we show that the cohomology groups $H^*(G(k,n),\mathbf{R})$ are generated by the first Pontrjagin class $p_1(E)$ and the Euler classes $e(E(k,n))$, $e(F(k,n))$ if k or $n-k$ are even. In these cases, the Poincaré duality: $H^q(G(k,n),\mathbf{R}) \to H_{k(n-k)-q}(G(k,n),\mathbf{R})$ can be given explicitly.

In §2, we compute volumes of some homogenous spaces which are needed in later discussion. In §3, we study the Poincaré duality on oriented compact Riemannian manifolds. The results are

Theorem 3.1 Let $\varphi_1, \cdots, \varphi_k$ be a basis of the cohomology group $H^q(M)$ represented by harmonic forms. Then $(\psi_1, \cdots, \psi_k) = (*\varphi_1, \cdots, *\varphi_k)A^{-1}$ is a basis of $H^{n-q}(M)$, where $a_{ij} = (\varphi_i, \varphi_j)$. The Poincaré duality D: $H^q(M) \to H_{n-q}(M)$ is given by

$$D(\varphi_i) = [T_i],$$

where $[T_1], \cdots, [T_k] \in H_{n-q}(M,\mathbf{R})$ are the dual of $\psi_1, \cdots, \psi_k$. Furthermore, if $[S_1], \cdots, [S_k]$ are the dual basis of $\varphi_1, \cdots, \varphi_k$, then

$$D(\psi_i) = (-1)^{q(n-q)}[S_i].$$

The Poincaré polynomials of Grassmann manifold $G(k,n)$ for $k=2$ and $n \leq 8$ are listed in §3, which give the structure of real homology and cohomology groups of Grassmann manifold. In §4, we show that the tangent space of Grassmann manifold is isomorphic to tensor product of the canonical vector bundles. Then we use the splitting principle of the characteristic class to study the relationship among these vector bundles, show that the characteristic classes of the tangent bundle on Grassmann manifold can be represented by that of canonical vector bundles.

In §5, we study $G(2,N)$, the main result are the following

Theorem 5.5 (1) When $2k+2 < N$, $[\mathbf{C}P^k]$ and $e^k(E(2,N))$ are the generators of $H_{2k}(G(2,N),\mathbf{Z})$ and $H^{2k}(G(2,N),\mathbf{Z})$ respectively;

(2) When $2k+2 > N$, $[G(2,k+2)]$ and $\frac{1}{2}e^k(E(2,N))$ are the generators of $H_{2k}(G(2,N),\mathbf{Z})$ and $H^{2k}(G(2,N),\mathbf{Z})$ respectively;

(3) When $2k+2 < N$, $D(e^k(E(2,N))) = [G(2,N-k)]$; when $2k+2 > N$, $D(\frac{1}{2}e^k(E(2,N))) = (-1)^{n-k}[\mathbf{C}P^{n-k-2}]$;

(4) $[\mathbf{C}P^n], [\overline{\mathbf{C}P}^n]$ and $\frac{1}{2}\{(-1)^n e^n(E(2,2n+2)) \pm e(F(2,2n+2))\}$ are generators of $H_{2n}(G(2,2n+2),\mathbf{Z})$ and $H^{2n}(G(2,2n+2),\mathbf{Z})$ respectively. Furthermore,

$$D(\frac{1}{2}[(-1)^n e^n(E(2,2n+2)) + e(F(2,2n+2))]) = [\mathbf{C}P^n],$$

$$D(\frac{1}{2}[(-1)^n e^n(E(2,2n+2)) - e(F(2,2n+2))]) = [\overline{\mathbf{C}P}^n].$$

In §6, we study the Grassmann manifold $G(3,6)$, the main result are

Theorem 6.1 (1) $\frac{1}{2}p_1(E(3,6)) \in H^4(G(3,6),\mathbf{Z})$ is a generator and its Poincaré dual $[SLAG]$ is a generator of $H_5(G(3,6),\mathbf{Z})$;

(2) $\frac{4}{3\pi} * p_1(E(3,6)) \in H^5(G(3,6),\mathbf{Z})$ is a generator and its Poincaré dual $[G(2,4)]$ is a generator of $H_4(G(3,6),\mathbf{Z})$.

In §7, §8 we study the Grassmann manifold $G(3,7)$ and $G(3,8)$, the main results are Theorem 7.5 and 8.4. In §9, we study $G(4,8)$, the main results are

Theorem 9.4 (1) $e(E), e(F), \frac{1}{2}(p_1(E)+e(E)-e(F)) \in H^4(G(4,8),\mathbf{Z})$ are the generators, their dual generators are $[\mathbf{C}P^2], [*\mathbf{C}P^2], [G(2,4)] \in H_4(G(4,8),\mathbf{Z})$;

(2) $\frac{1}{2}e^3(E), \frac{1}{2}e^3(F), \frac{1}{2}p_1^3(E)$ and $[G(4,7)], [G(3,7)], [CAY]$ are the generators of $H^{12}(G(4,8),\mathbf{Z})$ and $H_{12}(G(4,8),\mathbf{Z})$ respectively;

(3) The Poincaré duals of $e(E), e(F), \frac{1}{2}(p_1(E)+e(E)-e(F))$ are

$$[G(4,7)],\ [G(3,7)],\ [CAY]+[G(4,7)]-[G(3,7)]$$

respectively.

Theorem 9.5 The characteristic classes

$$\frac{1}{2}(e^2(F)+p_1(E)e(F)),\ \frac{1}{2}(e^2(F)-p_1(E)e(F)),$$

$$\frac{1}{2}(e^2(E)+p_1(E)e(E)),\ \frac{1}{2}(e^2(E)-p_1(E)e(E))$$

are the generators of $H^8(G(4,8),\mathbf{Z})$. Their Poincaré duals are

$$[ASSOC],\ [A\widetilde{SSO}C],\ [*ASSOC],\ [*A\widetilde{SSO}C]$$

respectively.

As application, in §5 and §9, we consider the Gauss maps of submanifolds in Euclidean spaces. For example, if $g\colon M \to G(4,8)$ is the Gauss map of an immersion $f\colon M \to \mathbf{R}^8$ of a compact oriented 4 dimensional manifold, we have

$$g_*[M] = \frac{1}{2}\chi(M)[G(4,5)] + \lambda[G(1,5)] + \frac{3}{2}Sign(M)[G(2,4)],$$

where $\lambda = \frac{1}{2}\int_M \ e(F(4,8))$ and $Sign(M)$ be the Signature of M. $\lambda = 0$ if f is an imbedding.

In §10 we use Gysin sequence to compute the cohomology of the homogenous space $ASSOC = G_2/SO(4)$, which had been studied by Borel and Hirzebruch [4].

The structure of the cohomology groups of infinite Grassmann manifold $G(k, \mathbf{R}^\infty)$ are simple, they are generated by Pontrjagin classes and the Euler class (if k is even) of the canonical vector bundle freely, see [12], p.179.

§2. The volumes of homogenous spaces

Let $e_1, e_2, \cdots, e_n$ be orthonormal frame fields on $\mathbf{R}^n$ such that $G(k,n)$ be generated by $e_1 \wedge \cdots \wedge e_k$ locally. The vectors $e_1, e_2, \cdots, e_n$ can be viewed as functions on Grassmann manifold. Let $\mathrm{d}e_A = \sum\limits_{B=1}^{n} \ \omega_A^B e_B, \ \ \omega_A^B = \langle \mathrm{d}e_A, e_B \rangle$ be 1 forms on $G(k,n)$. From $\mathrm{d}^2 e_A = 0$, we have $\mathrm{d}\omega_A^B = \sum\limits_{C=1}^{n} \ \omega_A^C \wedge \omega_C^B$. By

$$\mathrm{d}(e_1 \wedge \cdots \wedge e_k) = \sum_{i=1}^{k} \sum_{\alpha=k+1}^{n} \omega_i^\alpha E_{i\alpha},$$

$$E_{i\alpha} = e_1 \cdots e_{i-1} e_\alpha e_{i+1} \cdots e_k, \ i = 1, \cdots, k, \ \alpha = k+1, \cdots, n,$$

we know $E_{i\alpha}$ form a basis of $T_{e_1 \cdots e_k} G(k,n)$, ω_i^α be their dual basis.

$$\mathrm{d}s^2 = \langle \mathrm{d}(e_1 \wedge \cdots \wedge e_k), \mathrm{d}(e_1 \wedge \cdots \wedge e_k) \rangle = \sum_{i=1}^{k} \sum_{\alpha=k+1}^{n} (\omega_i^\alpha)^2$$

is the induced metric on $G(k,n)$. Differential $E_{i\alpha} = e_1 \cdots e_{i-1} e_\alpha e_{i+1} \cdots e_k$, by Gauss equation, we get the Riemannian connection ∇ on $G(k,n)$,

$$\nabla E_{i\alpha} = \sum_{j=1}^{k} \ \omega_i^j E_{j\alpha} + \sum_{\beta=k+1}^{n} \ \omega_\alpha^\beta E_{i\beta}.$$

Grassmann manifold $G(k,n)$ is oriented, the orientation is given by the volume form

$$\omega_1^{k+1} \wedge \omega_2^{k+1} \wedge \cdots \wedge \omega_k^{k+1} \wedge \cdots \wedge \omega_1^n \wedge \omega_2^n \wedge \cdots \wedge \omega_k^n.$$

For later use we compute the volume of some homogenous spaces, such as real Grassmann manifold $G(k,n)$, complex Grassmann manifold $G_{\mathbf{C}}(k,n)$ and $SLAG = SU(n)/SO(n)$. We first compute the volume of special orthogonal group $SO(n)$.

Let $gl(n,\mathbf{R})$ be the set of all $n\times n$ real matrices with the inner product

$$\langle X,Y\rangle=\mathrm{tr}\,(XY^t)=\sum_{A,B}X_{AB}Y_{AB},\quad X=(X_{AB}),Y=(Y_{CD})\in gl(n,\mathbf{R}).$$

Then $gl(n,\mathbf{R})$ is an Euclidean space and $SO(n)$ is a Riemannian submanifold of $gl(n,\mathbf{R})$. Represent the elements of $SO(n)$ by $G=(e_1,\cdots,e_n)^t$, where e_A the A-th row of G. The vectors $e_1,\cdots,e_n$ can be viewed as functions of $SO(n)$, then $\omega_A^B=\langle \mathrm{d}e_A,e_B\rangle=\mathrm{d}e_A\cdot e_B^t$ are 1 forms on $SO(n)$, $\omega_A^B+\omega_A^B=0$. Let E_{BC} be the matrix with 1 in the B-th row, C-th column, the others being zero. We have

$$\mathrm{d}GG^{-1}=(\omega_A^B)=\sum_{A,B}\omega_A^BE_{AB},\quad \mathrm{d}G=\sum_{A,B}\omega_A^BE_{AB}G=\sum_{A<B}\omega_A^B(E_{AB}-E_{BA})G.$$

Then $\{(E_{AB}-E_{BA})G\}$ is a basis of $T_GSO(n)$ and

$$\mathrm{d}s^2=\langle \mathrm{d}G,\mathrm{d}G\rangle=2\sum_{A<B}\omega_A^B\otimes\omega_A^B$$

is Riemannian metric on $SO(n)$.

Proposition 2.1 The volume of $SO(n)$ is

$$V(SO(n))=2^{\frac{1}{2}(n-1)}V(S^{n-1})V(SO(n-1))=2^{\frac{1}{4}n(n-1)}V(S^{n-1})\cdots V(S^1).$$

Proof Let $\bar{e}_n=(0,\cdots,0,1)$ be a fixed vector. The map $\tau(G)=\bar{e}_nG=e_n$ defines a fibre bundle τ: $SO(n)\to S^{n-1}$ with fibres $SO(n-1)$. By $\mathrm{d}e_n=\sum\omega_n^Ae_A$,

$$\mathrm{d}V_{S^{n-1}}=\omega_1^n\cdots\omega_{n-1}^n$$

is the volume element of S^{n-1}. The volume element of $SO(n)$ can be represented by

$$\mathrm{d}V_{SO(n)}=(\sqrt{2})^{\frac{1}{2}n(n-1)}\prod_{A<B}\omega_A^B=2^{\frac{1}{2}(n-1)}(\sqrt{2})^{\frac{1}{2}(n-1)(n-2)}\prod_{A<B<n}\omega_A^B\cdot\tau^*\mathrm{d}V_{S^{n-1}},$$

restricting $(\sqrt{2})^{\frac{1}{2}(n-1)(n-2)}\prod_{A<B<n}\omega_A^B$ on the fibres of τ are the volume elements of the fibres. Integration $\mathrm{d}V_{SO(n)}$ along the fibre of τ first, then on S^{n-1}, these shows

$$V(SO(n))=2^{\frac{1}{2}(n-1)}V(S^{n-1})V(SO(n-1)).\quad\square$$

As we know $V(S^m)=\frac{2\pi^{\frac{m+1}{2}}}{\Gamma(\frac{m+1}{2})}$,

$$V(S^{2n-1})=\frac{2\pi^n}{(n-1)!},\quad V(S^{2n})=\frac{2^{2n+1}n!\pi^n}{(2n)!}.$$

To compute the volume of $G(k,n)$, we use principle bundle $SO(n) \to G(k,n)$ with the Lie group $SO(k) \times SO(n-k)$ as fibres. The proof is similar to that of Proposition 2.1.

Proposition 2.2 The volume of Grassmann manifold $G(k,n)$ is

$$V(G(k,n)) = \frac{V(SO(n))}{2^{\frac{1}{2}k(n-k)}V(SO(k))V(SO(n-k))} = \frac{V(S^{n-1})\cdots V(S^{n-k})}{V(S^{k-1})\cdots V(S^1)}.$$

By simple computation, we have

$$V(G(2,n+2)) = \frac{2(2\pi)^n}{n!}, \quad V(G(3,6)) = \frac{2}{3}\pi^5,$$

$$V(G(3,7)) = \frac{16}{45}\pi^6, \quad V(G(3,8)) = \frac{2}{45}\pi^8, \quad V(G(4,8)) = \frac{8}{135}\pi^8.$$

Now we compute the volume of complex Grassmann manifold $G_{\mathbf{C}}(k,n)$. Let J be the natural complex structure on $\mathbf{C}^n = \mathbf{R}^{2n}$ and $s_1, \cdots, s_k$ be Hermitian orthonormal basis of $\pi \in G_{\mathbf{C}}(k,n)$. Let e_{2i-1}, $e_{2i} = Je_{2i-1} \in \mathbf{R}^{2n}$ be the realization vectors of s_i, $\sqrt{-1}\,s_i$ respectively. Then $e_1e_2\cdots e_{2k-1}e_{2k} \in G(2k,2n)$, and $G_{\mathbf{C}}(k,n)$ becomes a submanifold of $G(2k,2n)$.

Let $U(n) = \{G \in gl(n,\mathbf{C}) \mid G\overline{G}^t = I\}$ be the unitary group, the Hermitian inner product of $X = (X_{AB}), Y = (Y_{CD}) \in gl(n,\mathbf{C})$ be

$$\langle X, Y\rangle = \mathrm{tr}\,(X\overline{Y}^t) = \sum_{A,B} X_{AB}\overline{Y}_{AB}.$$

Let $G = (s_1, \cdots, s_n)^t \in U(n)$ represented by the rows of G, $\omega_A^B = \langle \mathrm{d}s_A, s_B\rangle = \mathrm{d}s_A \cdot \bar{s}_B^t$ be 1 forms on $U(n)$. Let $\omega_A^B = \varphi_A^B + \sqrt{-1}\,\psi_A^B$. From $\omega_A^B + \overline{\omega}_B^A = 0$ we have $\varphi_A^B + \varphi_B^A = 0$, $\psi_A^B - \psi_B^A = 0$. Then

$$\begin{aligned}\mathrm{d}G &= \sum_{A,B} \omega_A^B E_{AB} G \\ &= \sum_{A<B} \varphi_A^B (E_{AB} - E_{BA})G + \sqrt{-1}\,\{\sum_{A<B} \psi_A^B (E_{AB} + E_{BA})G + \sum_A \psi_A^A E_{AA} G\},\end{aligned}$$

and

$$\mathrm{d}s^2 = \langle \mathrm{d}G, \mathrm{d}G\rangle = 2\sum_{A<B}(\varphi_A^B \otimes \varphi_A^B + \psi_A^B \otimes \psi_A^B) + \sum_A \psi_A^A \otimes \psi_A^A$$

is Riemannian metric on $U(n)$. The volume element is

$$\mathrm{d}V_{U(n)} = 2^{\frac{1}{2}n(n-1)}\psi_1^1 \cdots \psi_n^n \prod_{A<B} \varphi_A^B \psi_A^B.$$

Proposition 2.3 (1) The volume of $U(n)$ is

$$V(U(n)) = 2^{n-1}V(S^{2n-1})V(U(n-1)) = 2^{\frac{1}{2}n(n-1)}V(S^{2n-1})V(S^{2n-3})\cdots V(S^1);$$

(2) As Riemannian submanifold of $G(2k,2n)$, the volume of $G_{\mathbf{C}}(k,n)$ is

$$V(G_{\mathbf{C}}(k,n)) = \frac{V(U(n))}{V(U(k))V(U(n-k))}.$$

Proof Let $\bar{e}_n = (0,\cdots,0,1)$ be a fixed vector. The map $\tau(G) = \bar{e}_n G = s_n$ defines a fibre bundle τ: $U(n) \to S^{2n-1}$ with fibre type $U(n-1)$. From $\mathrm{d}s_n = \sum \omega_n^A s_A$ and $\omega_n^A = \varphi_n^A + \sqrt{-1}\,\psi_n^A$, $\varphi_n^n = 0$, we have the volume element of S^{2n-1},

$$\mathrm{d}V_{S^{2n-1}} = \varphi_1^n\psi_1^n\cdots\varphi_{n-1}^n\psi_{n-1}^n\psi_n^n.$$

Then the volume element of $U(n)$ can be represented by

$$\mathrm{d}V_{U(n)} = 2^{n-1}\tau^*\mathrm{d}V_{S^{2n-1}}\cdot \mathrm{d}V_{U(n-1)}.$$

These prove (1).

As noted above, the map $[s_1\cdots s_k] \mapsto e_1e_2\cdots e_{2k-1}e_{2k}$ gives an imbedding of $G_{\mathbf{C}}(k,n)$ in $G(2k,2n)$. From $\mathrm{d}s_i = \sum\limits_{j=1}^{k} \omega_i^j s_j + \sum\limits_{\alpha=k+1}^{n} \omega_i^\alpha s_\alpha$, $\omega_i^j = \varphi_i^j + \sqrt{-1}\,\psi_i^j$, $\omega_i^\alpha = \varphi_i^\alpha + \sqrt{-1}\,\psi_i^\alpha$, we have

$$\mathrm{d}e_{2i-1} = \sum_{j=1}^{k} (\varphi_i^j e_{2j-1} + \psi_i^j e_{2j}) + \sum_{\alpha=k+1}^{n} (\varphi_i^\alpha e_{2\alpha-1} + \psi_i^\alpha e_{2\alpha}),$$

$$\mathrm{d}e_{2i} = \sum_{j=1}^{k} (\varphi_i^j e_{2j} - \psi_i^j e_{2j-1}) + \sum_{\alpha=k+1}^{n} (\varphi_i^\alpha e_{2\alpha} - \psi_i^\alpha e_{2\alpha-1}).$$

Then

$$\mathrm{d}(e_1e_2\cdots e_{2k-1}e_{2k}) = \sum_{i,\alpha} \varphi_i^\alpha (E_{2i-1\,2\alpha-1} + E_{2i\,2\alpha}) + \sum_{i,\alpha} \psi_i^\alpha (E_{2i-1\,2\alpha} - E_{2i\,2\alpha-1}),$$

$$\mathrm{d}V_{G_{\mathbf{C}}(k,n)} = 2^{k(n-k)}\varphi_1^{k+1}\psi_1^{k+1}\cdots\varphi_k^n\psi_k^n.$$

The rest is similar to that of Proposition 2.1. $\square$

The symmetric space $SLAG = SU(n)/SO(n)$ can be imbedded in $G(n,2n)$ as follows. Let $\bar{e}_{2i-1}, \bar{e}_{2i} = J\bar{e}_{2i-1}$, $i = 1,\cdots,n$, be a fixed orthonormal basis of $\mathbf{C}^n = \mathbf{R}^{2n}$, $M = \{G(\bar{e}_1\bar{e}_3\cdots\bar{e}_{2n-1}) \mid G \in SU(n) \subset SO(2n)\}$ is diffeomorphic to $SLAG = SU(n)/SO(n)$.

Proposition 2.4 (1) The volume of special unitary group $SU(n)$ is

$$V(SU(n)) = 2^{n-1}\sqrt{\frac{n}{n-1}}V(S^{2n-1})V(SU(n-1));$$

(2) The volume of $M = SLAG$ is

$$V(M) = \frac{V(SU(n))}{V(SO(n))}.$$

Proof The proof is similar to Proposition 2.1. Let $G = (s_1, \cdots, s_n)^t \in SU(n)$, $\omega_A^B = \mathrm{d}s_A \cdot \bar{s}_B^t$. From $\det G = 1$ we have $\sum\limits_{A=1}^{n} \omega_A^A = 0$, then $\psi_n^n = -\sum\limits_{B\neq n} \psi_B^B$. The Riemannian metric on $SU(n)$ is

$$\begin{aligned}
\mathrm{d}s^2 &= 2\sum_{A<B}(\varphi_A^B \otimes \varphi_A^B + \psi_A^B \otimes \psi_A^B) + \sum_{B\neq n}\psi_B^B \otimes \psi_B^B + \psi_n^n \otimes \psi_n^n \\
&= 2\sum_{A<B}(\varphi_A^B \otimes \varphi_A^B + \psi_A^B \otimes \psi_A^B) \\
&\quad +(\psi_1^1, \cdots, \psi_{n-1}^{n-1})\begin{pmatrix} 2 & 1 & \cdots & 1 \\ 1 & 2 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 2 \end{pmatrix}\begin{pmatrix} \psi_1^1 \\ \vdots \\ \psi_{n-1}^{n-1} \end{pmatrix}.
\end{aligned}$$

Then

$$\mathrm{d}V_{SU(n)} = 2^{\frac{1}{2}n(n-1)}\sqrt{n}\psi_1^1 \cdots \psi_{n-1}^{n-1} \prod_{A<B} \varphi_A^B \psi_A^B.$$

The volume of special unitary group $SU(n)$ is

$$V(SU(n)) = 2^{n-1}\sqrt{\frac{n}{n-1}}V(S^{2n-1})V(SU(n-1)).$$

Let $e_{2A-1},\ e_{2A} = Je_{2A-1}$ be the realization vectors of $s_A,\ \sqrt{-1}\,s_A$ respectively. $M = SLAG$ is generated by $G(\bar{e}_1\bar{e}_3 \cdots \bar{e}_{2n-1}) = e_1 e_3 \cdots e_{2n-1}$,

$$\mathrm{d}(e_1 e_3 \cdots e_{2n-1}) = \sum \psi_B^B(E_{2B-1\,2B} - E_{2n-1\,2n}) + \sum_{A<B} \psi_A^B(E_{2A-1\,2B} + E_{2B-1\,2A}),$$

$$\mathrm{d}s^2 = 2\sum_{A<B}\psi_A^B \otimes \psi_A^B + 2\sum_{B\neq n}\psi_B^B \otimes \psi_B^B + \sum_{B\neq C<n}\psi_B^B \otimes \psi_C^C.$$

Then

$$\mathrm{d}V_M = 2^{\frac{1}{4}n(n-1)}\sqrt{n}\psi_1^1 \cdots \psi_{n-1}^{n-1} \prod_{A<B} \psi_A^B.$$

Let τ: $SU(n) \to M$ be the projection, the fibre of τ at $e_1 e_3 \cdots e_{2n-1}$ is the group $SO(n)$. Restricting $\mathrm{d}s_i = \sum\limits_{j=1}^{k} \omega_i^j s_j + \sum\limits_{\alpha=k+1}^{n} \omega_i^\alpha s_\alpha$ on the fibre of τ, we have $\omega_i^\alpha = 0$ and $\psi_i^j = 0$, then $\mathrm{d}V_{SO(n)} = 2^{\frac{1}{4}n(n-1)} \prod_{A<B} \varphi_A^B$ is the volume element of the fibres. These completes the proof. $\square$

§3. The Poincaré duality

Let M be a compact oriented Riemannian manifold and $H_q(M) = H_q(M, \mathbf{R})$ its q-th singular homology group, $H^q(M) = H^q(M, \mathbf{R})$ be the q-th de Rham cohomology group. For any $[\xi] \in H^q(M)$ and $[z] = [\sum \lambda_i \sigma_i] \in H_q(M)$, we can define

$$[\xi]([z]) = \int_z \xi = \sum \lambda_i \int_{\sigma_i} \xi = \sum \lambda_i \int_{\triangle^q} \sigma_i^* \xi,$$

where every singular simplex σ_i: $\triangle^q \to M$ is differentiable. If $[\xi] \in H^q(M, \mathbf{Z})$ and $[z] \in H_q(M, \mathbf{Z})$, we have $[\xi]([z]) \in \mathbf{Z}$. By universal coefficients Theorem, we have

$$H^q(M, \mathbf{R}) \cong \mathrm{Hom}\,(H_q(M, \mathbf{R}), \mathbf{R}),$$

and

$$H^q(M, \mathbf{Z}) \cong \mathrm{Hom}\,(H_q(M, \mathbf{Z}), \mathbf{Z}) \oplus \mathrm{Ext}\,(H_{q-1}(M, \mathbf{Z}), \mathbf{Z}).$$

On the other hand, we have Poincaré duality

$$D\colon\ H^q(M, \mathbf{R}) \to H_{n-q}(M, \mathbf{R}), \quad n = \dim M.$$

For any $[\xi] \in H^q(M)$, $D[\xi] \in H_{n-q}(M)$, we have

$$[\eta](D[\xi]) = \int_{D[\xi]} \eta = \int_M \xi \wedge \eta$$

for any $[\eta] \in H^{n-q}(M)$.

In the following, we use harmonic forms to represent the Poincaré duality. Let $\varphi_1, \cdots, \varphi_k$ be the basis of $H^q(M)$ and $[T_i] = D(\varphi_i)$ be their Poincaré duals. By Hodge Theorem, we can assume that $\varphi_1, \cdots, \varphi_k$ are all the harmonic forms on M. Then $*\varphi_1, \cdots, *\varphi_k$ are the harmonic forms on M and form a basis of $H^{n-q}(M)$. Let

$$a_{ij} = (\varphi_i, \varphi_j) = \int_M \langle \varphi_i, \varphi_j \rangle \, \mathrm{d}V_M = \int_M \varphi_i \wedge *\varphi_j$$

be the inner product of differential forms φ_i, φ_j. Let $\psi_1, \cdots, \psi_k$ be the dual basis of $[T_1], \cdots, [T_k]$, also represented by harmonic forms. Assuming $\psi_j = \sum *\varphi_i b_{ij}$, by Poincaré duality,

$$\delta_{ij} = \int_{T_i} \psi_j = \int_M \varphi_i \wedge \psi_j = \int_M \sum \varphi_i \wedge *\varphi_l b_{lj} = \sum a_{il} b_{lj}.$$

This shows $(b_{ij}) = (a_{ij})^{-1} = A^{-1}$, we have

$$(\psi_1, \cdots, \psi_k) = (*\varphi_1, \cdots, *\varphi_k)A^{-1}.$$

Theorem 3.1 Let $\varphi_1, \cdots, \varphi_k$ be a basis of the cohomology group $H^q(M)$ represented by harmonic forms. Then $(\psi_1, \cdots, \psi_k) = (*\varphi_1, \cdots, *\varphi_k)A^{-1}$ is a basis of $H^{n-q}(M)$, where $a_{ij} = (\varphi_i, \varphi_j)$. The Poincaré duality D: $H^q(M) \to H_{n-q}(M)$ is given by

$$D(\varphi_i) = [T_i],$$

where $[T_1], \cdots, [T_k] \in H_{n-q}(M, \mathbf{R})$ are the dual of $\psi_1, \cdots, \psi_k$. Furthermore, if $[S_1], \cdots, [S_k]$ are the dual basis of $\varphi_1, \cdots, \varphi_k$, then

$$D(\psi_i) = (-1)^{q(n-q)}[S_i].$$

Proof The equations $D(\psi_i) = (-1)^{q(n-q)}[S_i]$ follow from $**\varphi_i = (-1)^{q(n-q)}\varphi_i$ and $(\varphi_i, \varphi_j) = (*\varphi_i, *\varphi_j)$. $\square$

Theorem 3.1 can be applied to the Poincaré duality D: $H^q(M, \mathbf{Z}) \to H_{n-q}(M, \mathbf{Z})$ if we ignore the torsion elements of $H^q(M, \mathbf{Z})$.

The q-th Betti number is the common dimension of the real homology and cohomology groups $H_q(G(k, n))$ and $H^q(G(k, n))$ (and is also the rank of $H_q(G(k, n), \mathbf{Z})$ and $H^q(G(k, n), \mathbf{Z})$). The Poincaré polynomials, with the Betti numbers as coefficients are given by the following Table (see[7], [8]).

Grassmannian	Poincaré polynomial
$G(1, n+1)$	$1 + t^n$
$G(2, 2n+1)$	$1 + t^2 + t^4 + \cdots + t^{4n-2}$
$G(2, 2n+2)$	$(1 + t^{2n})(1 + t^2 + \cdots + t^{2n})$
$G(3, 6)$	$(1 + t^4)(1 + t^5)$
$G(3, 7)$	$(1 + t^4 + t^8)(1 + t^4)$
$G(3, 8)$	$(1 + t^4 + t^8)(1 + t^7)$
$G(4, 8)$	$(1 + t^4 + t^8)(1 + t^4)^2$

§4. The vector bundles on $G(k, n)$

Let τ_1: $E(k, n) \to G(k, n)$ be the canonical vector bundle on Grassmann manifold $G(k, n)$, the fibre over $\pi \in G(k, n)$ be the vectors of π. $E = E(k, n)$ is a Riemannian vector bundle with the induced metric. Let $e_1, \cdots, e_k, e_{k+1}, \cdots, e_n$ be orthonormal frame fields on $\mathbf{R}^n$, $G(k, n)$ is locally generated by $e_1 \cdots e_k = e_1 \wedge \cdots \wedge e_k$.

Then $e_1, \cdots, e_k$ are local orthonormal sections of the vector bundle τ_1. From $\mathrm{d}e_i = \sum\limits_{j=1}^{k} \omega_i^j e_j + \sum\limits_{\alpha=k+1}^{n} \omega_i^\alpha e_\alpha$, we know that $\nabla e_i = \sum\limits_{j=1}^{k} \omega_i^j e_j$ defines a Riemannian connection on τ_1. From $\nabla^2 e_i = \sum\limits_{j=1}^{k} (\mathrm{d}\omega_i^j - \sum\limits_{l=1}^{k} \omega_i^l \wedge \omega_l^j) e_j$, we have curvature forms

$$\Omega_i^j = \mathrm{d}\omega_i^j - \sum_{l=1}^{k} \omega_i^l \wedge \omega_l^j = \sum_{\alpha=k+1}^{n} \omega_i^\alpha \wedge \omega_\alpha^j.$$

The total Pontrjagin classes of the vector bundle τ_1: $E \to G(k,n)$ are

$$p(E) = 1 + p_1(E) + p_2(E) + \cdots = \det(I + \frac{1}{2\pi}(\Omega_i^j)).$$

If k is even, we have Euler class,

$$e(E) = \frac{(-1)^{\frac{k}{2}}}{(4\pi)^{\frac{k}{2}}(\frac{k}{2})!} \sum_{i_1,\cdots,i_k} \varepsilon(i_1 i_2 \cdots i_k) \Omega_{i_1 i_2} \Omega_{i_3 i_4} \cdots \Omega_{i_{k-1} i_k}.$$

Similarly, we can define vector bundle τ_2: $F = F(k,n) \to G(k,n)$ on Grassmann manifold $G(k,n)$, the fibre over $e_1 \cdots e_k \in G(k,n)$ is the vectors orthogonal to $e_1, \cdots, e_k$. Then $e_{k+1}, \cdots, e_n$ are local orthonormal sections of F. From $\mathrm{d}e_\alpha = \sum\limits_{\beta=k+1}^{n} \omega_\alpha^\beta e_\beta + \sum\limits_{i=1}^{k} \omega_\alpha^i e_i$, we have Riemannian connection $\nabla e_\alpha = \sum\limits_{\beta=k+1}^{n} \omega_\alpha^\beta e_\beta$. The curvature forms are given by

$$\nabla^2 e_\alpha = \sum_{j=1}^{k} \Omega_\alpha^\beta e_\beta, \quad \Omega_\alpha^\beta = \sum_{i=1}^{k} \omega_\alpha^i \wedge \omega_i^\beta.$$

The total Pontrjagin classes of the vector bundle τ_2: $F \to G(k,n)$ are

$$p(F) = 1 + p_1(F) + p_2(F) + \cdots = \det(I + \frac{1}{2\pi}(\Omega_\alpha^\beta)).$$

The direct sum $E(k,n) \oplus F(k,n) = G(k,n) \times \mathbf{R}^n$ is trivial, we have

$$(1 + p_1(E) + p_2(E) + \cdots) \cdot (1 + p_1(F) + p_2(F) + \cdots) = 1.$$

Then Pontrjagin classes of F are determined by that of E. For example, we have

$$p_1(F) = -p_1(E), \ \ p_2(F) = p_1^2(E) - p_2(E), \ \ p_3(F) = -p_1^3(E) + 2p_1(E)p_2(E) - p_3(E).$$

Let $*$: $\bigwedge^k(\mathbf{R}^n) \to \bigwedge^{n-k}(\mathbf{R}^n)$ be the star operator, $*G(k,n) = G(n-k,n)$ and the canonical vector bundles $E(k,n), F(k,n)$ are interchanged under the map $*$.

Proposition 4.1 The tangent space $TG(k,n)$ of Grassmann manifold is isomorphic to tensor product $E(k,n)\otimes F(k,n)$. If $k(n-k)$ is even, we have

$$e(G(k,n)) = e(E(k,n)\otimes F(k,n)).$$

Proof Let $e_1, e_2, \cdots, e_n$ be an oriented orthonormal basis of $\mathbf{R}^n$, the fibre of $E(k,n)$ over $x = e_1\wedge\cdots\wedge e_k \in G(k,n)$ is generated by $e_1,\cdots,e_k$ and the fibre of $F(k,n)$ over x is generated by $e_{k+1},\cdots,e_n$. On the other hand, the tangent space $T_xG(k,n)$ is generated by $E_{i\,\alpha} = e_1\wedge\cdots\wedge e_{i-1}\wedge e_\alpha\wedge e_{i+1}\cdots\wedge e_k$. It is easy to see that the map $E_{i\,\alpha}\mapsto e_i\otimes e_\alpha$ gives an isomorphism from tangent bundle $TG(k,n)$ to tensor product $E\otimes F$.

It is easy to see that the isomorphism $TG(k,n)\to E(k,n)\otimes F(k,n)$ preserves the connections on $TG(k,n)$ and $E\otimes F$ respectively, where the connection on $E\otimes F$ is

$$\nabla\,(e_i\otimes e_\alpha) = \sum_{j=1}^{k}\,\omega_i^j e_j\otimes e_\alpha + \sum_{\alpha=k+1}^{n}\,\omega_\alpha^\beta e_i\otimes e_\beta. \quad \square$$

In the following we use the splitting principle of the characteristic class to study the relationship among these vector bundles. First we study the oriented Grassmann manifold $G(2k,2n)$. Assuming $s_1,\cdots,s_{2k}$ be the orthonormal sections of vector bundle $E(2k,2n)$ such that the curvature of Riemannian connection has the form

$$\frac{1}{2\pi}\nabla^2\begin{pmatrix} s_1\\ s_2\\ \vdots\\ s_{2k-1}\\ s_{2k}\end{pmatrix} = \begin{pmatrix} 0 & -x_1 & & & \\ x_1 & 0 & & & \\ & & \ddots & & \\ & & & 0 & -x_k\\ & & & x_k & 0\end{pmatrix}\begin{pmatrix} s_1\\ s_2\\ \vdots\\ s_{2k-1}\\ s_{2k}\end{pmatrix}.$$

The total Pontrjagin classes and the Euler class of $E = E(2k,2n)$ are

$$p(E) = \prod_{i=1}^{k}\,(1+x_i^2), \quad e(E) = x_1\cdots x_k.$$

Similarly assuming $t_{2k+1}, t_{2k+2},\cdots,t_{2n}$ be the orthonormal sections of vector bundle $F(2k,2n)$, the curvature of Riemannian connection has the form

$$\frac{1}{2\pi}\nabla^2\begin{pmatrix} t_{2k+1}\\ t_{2k+2}\\ \vdots\\ t_{2n-1}\\ t_{2n}\end{pmatrix} = \begin{pmatrix} 0 & -y_{k+1} & & & \\ y_{k+1} & 0 & & & \\ & & \ddots & & \\ & & & 0 & -y_n\\ & & & y_n & 0\end{pmatrix}\begin{pmatrix} t_{2k+1}\\ t_{2k+2}\\ \vdots\\ t_{2n-1}\\ t_{2n}\end{pmatrix}.$$

The total Pontrjagin classes and the Euler class of $F = F(2k, 2n)$ are

$$p(F) = \prod_{\alpha=k+1}^{n} (1 + y_\alpha^2), \quad e(F) = y_{k+1} \cdots y_n.$$

$s_{2i-1} \otimes t_{2\alpha-1}, s_{2i} \otimes t_{2\alpha-1}, s_{2i-1} \otimes t_{2\alpha}, s_{2i} \otimes t_{2\alpha}$ are the local orthonormal sections of vector bundle $E \otimes F \cong TG(2k, 2n)$. The curvature of Riemannian connection on $E \otimes F$ is given by

$$\frac{1}{2\pi}\nabla^2 \begin{pmatrix} s_{2i-1} \otimes t_{2\alpha-1} \\ s_{2i} \otimes t_{2\alpha-1} \\ s_{2i-1} \otimes t_{2\alpha} \\ s_{2i} \otimes t_{2\alpha} \end{pmatrix} = \begin{pmatrix} 0 & -x_i & -y_\alpha & 0 \\ x_i & 0 & 0 & -y_\alpha \\ y_\alpha & 0 & 0 & -x_i \\ 0 & y_\alpha & x_i & 0 \end{pmatrix} \begin{pmatrix} s_{2i-1} \otimes t_{2\alpha-1} \\ s_{2i} \otimes t_{2\alpha-1} \\ s_{2i-1} \otimes t_{2\alpha} \\ s_{2i} \otimes t_{2\alpha} \end{pmatrix}.$$

Then we have

Lemma 4.2 (1) $e(TG(2k, 2n)) = e(E \otimes F) = \prod_{i,\alpha} (x_i^2 - y_\alpha^2)$;

(2) $p(TG(2k, 2n)) = p(E \otimes F) = \prod_{i,\alpha} (1 + 2(x_i^2 + y_\alpha^2) + (x_i^2 - y_\alpha^2)^2)$.

The other cases can be discussed similarly.

By simple computation, we have

$$p_1(TG(2k, 2n)) = (2n - 2k)p_1(E) + 2kp_1(F) = 2(n - 2k)p_1(E).$$

In particular, $p_1(TG(2k, 4k)) = 0$.

In next section, we shall show

$$e(TG(2, 2n + 2)) = (n + 1)e^{2n}(E(2, 2n + 2)),$$

$$e(TG(2, 2n + 3)) = (n + 1)e^{2n+1}(E(2, 2n + 3)).$$

We can also show

$$e(TG(3, 7)) = 3e^3(F(3, 7)), \quad e(TG(4, 8)) = 6e^4(E(4, 8)) = 6e^4(F(4, 8)).$$

§5. The cases of $G(2, N)$

In this section, we study the real homology and cohomology of Grassmann manifold $G(2, N)$.

As is well-known, the oriented Grassmann manifold $G(2, N)$ is Kaehler manifold and can be imbedded in complex projective space. Here we give a new proof. Let

e_1, e_2 be oriented orthonormal basis of $\pi \in G(2,N)$, $e_1 \mapsto e_2$, $e_2 \mapsto -e_1$ defines an almost complex structure

$$J\colon\ T_\pi G(2,N) \to T_\pi G(2,N),$$

$$E_{1\,\alpha} = e_\alpha \wedge e_2 \mapsto -e_\alpha \wedge e_1 = E_{2\,\alpha},\ E_{2\,\alpha} = e_1 \wedge e_\alpha \mapsto e_2 \wedge e_\alpha = -E_{1\,\alpha}.$$

It is easy to see that J is well defined and preserves the metric on $G(2,N)$.

Proposition 5.1 $G(2,N)$ is a Kaehler manifold with complex structure J.

Proof Let ∇ be the Riemannian connection on $TG(2,N)$ defined above. We have

$$(\nabla J)E_{i\,\alpha} = \nabla(JE_{i\,\alpha}) - J(\nabla E_{i\,\alpha}) = 0, \quad i = 1, 2.$$

Hence, $\nabla J = 0$, J is a complex structure and $G(2,N)$ is a Kaehler manifold. □

The Euler classes of canonical vector bundles $E = E(2, 2n+2)$ and $F = F(2, 2n+2)$ can be represented by

$$e(E) = \frac{1}{2\pi}\sum_{\alpha=3}^{2n+2} \omega_1^\alpha \wedge \omega_2^\alpha,$$

$$e(F) = \frac{(-1)^n}{(4\pi)^n n!}\sum \varepsilon(\alpha_1\alpha_2\cdots\alpha_{2n})\Omega_{\alpha_1\alpha_2}\wedge\cdots\wedge\Omega_{\alpha_{2n-1}\alpha_{2n}}.$$

$G(2, k+2)$ is a submanifold of $G(2, 2n+2)$ whose elements contained in a fixed $k+2$ dimensional subspace of $\mathbf{R}^{2n+2}$, $i\colon\ G(2,k+2) \to G(2,2n+2)$ the inclusion. Then, $E(2,k+2) = i^*E(2,2n+2)$ and $e(E(2,k+2)) = i^*e(E(2,2n+2))$. Let $G(1, 2n+1)$ be a submanifold of $G(2, 2n+2)$ with elements $e_1 \wedge \bar{e}_2$, $\bar{e}_2 = (0,\cdots,0,1)$, $j\colon\ G(1,2n+1) \to G(2,2n+2)$ be the inclusion.

Theorem 5.2 For Grassmann manifold $G(2, 2n+2)$, we have

(1) $p_q(F) = (-1)^q p_1^q(E) = (-1)^q e^{2q}(E), \quad q = 1, \cdots, n;$

(2) The Pontrjagin classes and Euler class of tangent bundle $TG(2, 2n+2)$ can be represented by the Euler class of E,

$$p_1(G(2,2n+2)) = 2(n-1)e^2(E),\quad p_2(G(2,2n+2)) = (2n^2-5n+9)e^4(E),\ \cdots,$$

$$e(G(2,2n+2)) = (n+1)e^{2n}(E);$$

(3) $\int_{G(1,2n+1)} e(F) = 2, \quad \int_{G(2,k+2)} e^k(E) = 2, \quad k = 1, \cdots, 2n.$

Proof For Grassmann manifold $G(2, 2n+2)$, we have $p_1(E) = e^2(E)$, $p_n(F) = e^2(F)$. From $(1 + p_1(E)) \cdot (1 + p_1(F) + p_2(F) + \cdots + p_n(F)) = 1$, we have

$$1 + p_1(F) + p_2(F) + \cdots + p_n(F) = \frac{1}{1 + p_1(E)} = 1 + \sum_{q=1}^{n} (-1)^q p_1^q(E).$$

Hence, $p_q(F) = (-1)^q p_1^q(E) = (-1)^q e^{2q}(E)$, $p_n(F) = e^2(F) = (-1)^n e^{2n}(E)$. These proves (1).

By Lemma 4.2, note that $x_1 = e(E)$, the Euler class of $G(2, 2n+2)$ is

$$\begin{aligned} e(TG(2,2n+2)) &= (x_1^2 - y_2^2)(x_1^2 - y_3^2)\cdots(x_1^2 - y_{n+1}^2) \\ &= x_1^{2n} - x_1^{2n-2}p_1(F) + x_1^{2n-4}p_2(F) - \cdots + (-1)^n p_n(F) \\ &= (n+1)e^{2n}(E). \end{aligned}$$

By Gauss-Bonnet formula, we have

$$\chi(G(2,2n+2)) = \int_{G(2,2n+2)} e(G(2,2n+2)) = 2n+2.$$

From (1) and

$$p(G(2,2n+2)) = \prod_{\alpha=2}^{n+1} (1 + 2(x_1^2 + y_\alpha^2) + (x_1^2 - y_\alpha^2)^2),$$

we can prove (2).

Restricting the Euler class $e(E)$ on $G(2, k+2)$, we have

$$i^* e(E) = \frac{1}{2\pi}\sum_{\alpha=3}^{k+2} \omega_1^\alpha \wedge \omega_2^\alpha.$$

Then

$$i^* e^k(E) = \frac{k!}{(2\pi)^k}\omega_1^3 \wedge \omega_2^3 \wedge \cdots \wedge \omega_1^{k+2} \wedge \omega_2^{k+2} = \frac{k!}{(2\pi)^k}\mathrm{d}\, V_{G(2,k+2)},$$

where $\mathrm{d}\, V_{G(2,k+2)}$ is the volume element of $G(2, k+2)$. Then

$$\int_{G(2,k+2)} e^k(E(2,2n+2)) = 2, \quad k = 1, \cdots, 2n.$$

Restricting on $G(1, 2n+1)$, $\omega_2^\alpha = 0$, we have

$$e(F(1,2n+1)) = j^* e(F(2,2n+2)) = \frac{(2n)!}{2^{2n} n! \pi^n}\omega_1^3 \wedge \omega_1^4 \wedge \cdots \wedge \omega_1^{2n+2}.$$

It is easy to see that $e(F(1, 2n+1))$ is the Euler class of the tangent bundle of $S^{2n} = G(1, 2n+1)$ and $\omega_1^3 \wedge \omega_1^4 \wedge \cdots \wedge \omega_1^{2n+2}$ is the volume element. Hence by Gauss-Bonnet formula or by direct computation, we have

$$\int_{G(1,2n+1)} j^* e(F(2,2n+2)) = 2.$$

Furthermore, from $\omega_2^\alpha|_{G(1,2n+1)} = 0$ and $\Omega_{\alpha\beta}|_{G(2,n+2)} = 0$ for $\alpha, \beta > n+2$, we have

$$\int_{G(1,2n+1)} j^* e^n(E(2,2n+2)) = 0, \quad \int_{G(2,n+2)} i^* e(F(2,2n+2)) = 0.$$

By $p_1^k(E(2,2n+2)) = e^{2k}(E(2,2n+2))$, we have

$$\int_{G(2,2k+2)} p_1^k(E(2,2n+2)) = 2. \quad \square$$

The Poincaré polynomial of $G(2,2n+2)$ is

$$p_t(G(2,2n+2)) = 1 + t^2 + \cdots + t^{2n-2} + 2t^{2n} + t^{2n+2} + \cdots + t^{4n}.$$

By Theorem 5.2, we have

(1) For $k \neq n$, $e^k(E) \in H^{2k}(G(2,2n+2))$, $G(2,k+2) \in H_{2k}(G(2,2n+2))$ are the generators respectively;

(2) $e^n(E), e(F) \in H^{2n}(G(2,2n+2))$ and $G(2,n+2), G(1,2n+1) \in H_{2n}(G(2,2n+2))$ are the generators.

The characteristic classes $e^k(E), e(F)$ and the submanifolds $G(2,k+2), G(1,2n+1)$ are integral cohomology and homology classes respectively. But they need not be the generators of the integral cohomology and homology groups. For example, when $k \neq n$, from $\int_{G(2,k+2)} e^k(E) = 2$ we know that $[G(2,k+2)] \in H_{2k}(G(2,2n+2), \mathbf{Z})$, $e^k(E) \in H^{2k}(G(2,2n+2), \mathbf{Z})$ can not be generators simultaneously. Now we compute $\int_{\mathbf{C}P^k} i^* e^k(E(2,2n+2))$ and $\int_{\mathbf{C}P^n} i^* e(F(2,2n+2))$.

Let J be a complex structure on $\mathbf{R}^{2n+2}$, $\mathbf{R}^{2k+2} \subset \mathbf{R}^{2n+2}$ be invariant under the action of J and $\mathbf{C}P^k = \{e_1 J e_1 \mid e_1 \in S^{2k+1}\}$. Let $e_1, e_2 = Je_1, e_{2\alpha-1}, e_{2\alpha} = Je_{2\alpha-1}$, $\alpha = 2, 3, \cdots, k+1$, be local Hermitian orthonormal frame fields on $\mathbf{R}^{2k+2}$. By $\mathrm{d}e_2 = J\mathrm{d}e_1$ we have $\omega_1^{2\alpha-1} = \omega_2^{2\alpha}$, $\omega_1^{2\alpha} = -\omega_2^{2\alpha-1}$, then

$$\mathrm{d}(e_1 \wedge e_2) = \sum_{\alpha=2}^{k+1} \omega_1^{2\alpha-1}(E_{1\,2\alpha-1} + E_{2\,2\alpha}) + \sum_{\alpha=2}^{k+1} \omega_1^{2\alpha}(E_{1\,2\alpha} - E_{2\,2\alpha-1}).$$

The oriented volume element of $\mathbf{C}P^k$ is $\mathrm{d}V = 2^k \omega_1^3 \wedge \omega_1^4 \wedge \cdots \wedge \omega_1^{2k+2}$.

Let i: $\mathbf{C}P^k \to G(2,2n+2)$ be inclusion, we have

$$i^* e^k(E) = (-1)^k \frac{k!}{\pi^k} \omega_1^3 \wedge \omega_1^4 \wedge \cdots \wedge \omega_1^{2k+2}.$$

By Proposition 2.3,

$$\int_{\mathbf{C}P^k} i^* e^k(E) = (-1)^k.$$

J induces a complex structure on the induced bundle $i^*F \to \mathbf{C}P^n$. Let $F_{\mathbf{C}}$ be the complex vector bundle formed by the $(1,0)$-vectors of $i^*F \otimes \mathbf{C}$. By $i^*e(F) = c_n(F_{\mathbf{C}})$ (see [20]), we can show

$$\int_{\mathbf{C}P^n} i^*e(F) = \int_{\mathbf{C}P^n} c_n(F_{\mathbf{C}}) = 1,$$

see also Chern [2].

Let $\bar{J}$ be a complex structure on $\mathbf{R}^{2k+2}$, the orientation given by $\bar{J}$ is opposite to that of J. Let $\overline{\mathbf{C}P}^k = \{v \wedge \bar{J}v \mid v \in S^{2k+1}\}$ be the complex projective space. The orientation on the vector bundle $E(2,2n+2)|_{\overline{\mathbf{C}P}^k}$ are given by $v, \bar{J}v$, we have

$$\int_{\overline{\mathbf{C}P}^k} i^*e^k(E) = (-1)^k.$$

By $p_k(E(2,2n+2)) = e^{2k}(E(2,2n+2))$, we have

$$\int_{\mathbf{C}P^{2k}} i^*p_k(E(2,2n+2)) = 1.$$

Let $\widetilde{F}_{\mathbf{C}}$ be the complex vector bundle formed by the $(1,0)$-vectors of $F \otimes \mathbf{C}|_{\overline{\mathbf{C}P}^n}$. The orientation on realization vector bundle of $\widetilde{F}_{\mathbf{C}}$ given by $\bar{J}$ is opposite to that of $F|_{\overline{\mathbf{C}P}^n}$. Hence $e(F|_{\overline{\mathbf{C}P}^n}) = -c_n(\widetilde{F}_{\mathbf{C}})$ and we have

$$\int_{\overline{\mathbf{C}P}^n} e(F) = -\int_{\overline{\mathbf{C}P}^n} c_n(\widetilde{F}_{\mathbf{C}}) = -1.$$

These prove

Proposition 5.3 (1) When $k < n$, we have

$$[G(2,k+2)] = 2(-1)^k[\mathbf{C}P^k] \in H_{2k}(G(2,2n+2));$$

(2) In the homology group $H_{2n}(G(2,2n+2))$, we have

$$[G(2,n+2)] = (-1)^n([\mathbf{C}P^n] + [\overline{\mathbf{C}P}^n]),$$

$$[G(1,2n+1)] = [\mathbf{C}P^n] - [\overline{\mathbf{C}P}^n].$$

For Grassmann manifold $G(2,2n+3)$, by the splitting principle of the characteristic classes, we can assume that there are oriented orthonormal sections s_1, s_2 and $t_3, t_4, \cdots, t_{2n+2}, t_{2n+3}$ of vector bundle $E(2k,2n+3)$ and $F(2,2n+3)$ respectively, such that

$$\frac{1}{2\pi}\nabla^2 \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{pmatrix} 0 & -x \\ x & 0 \end{pmatrix} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix},$$

$$\frac{1}{2\pi}\nabla^2\begin{pmatrix} t_3 \\ t_4 \\ \vdots \\ t_{2n+1} \\ t_{2n+2} \\ t_{2n+3} \end{pmatrix} = \begin{pmatrix} 0 & -y_2 & & & & \\ y_2 & 0 & & & & \\ & & \ddots & & & \\ & & & 0 & -y_{n+1} & \\ & & & y_{n+1} & 0 & \\ & & & & & 0 \end{pmatrix}\begin{pmatrix} t_3 \\ t_4 \\ \vdots \\ t_{2n+1} \\ t_{2n+2} \\ t_{2n+3} \end{pmatrix}.$$

The total Pontrjagin classes of F are $p(F) = \prod_{\alpha=2}^{n+1}(1+y_\alpha^2)$.

$s_1 \otimes t_{2\alpha-1}, s_2 \otimes t_{2\alpha-1}, s_1 \otimes t_{2\alpha}, s_2 \otimes t_{2\alpha}$ and $s_1 \otimes t_{2n+3}, s_2 \otimes t_{2n+3}$ are orthonormal sections of $E \otimes F \cong TG(2k, 2n)$, they also give an orientation of $E \otimes F$. The curvature of $E \otimes F$ are defined by

$$\frac{1}{2\pi}\nabla^2\begin{pmatrix} s_1 \otimes t_{2\alpha-1} \\ s_2 \otimes t_{2\alpha-1} \\ s_1 \otimes t_{2\alpha} \\ s_2 \otimes t_{2\alpha} \end{pmatrix} = \begin{pmatrix} 0 & -x & -y_\alpha & 0 \\ x & 0 & 0 & -y_\alpha \\ y_\alpha & 0 & 0 & -x \\ 0 & y_\alpha & x & 0 \end{pmatrix}\begin{pmatrix} s_1 \otimes t_{2\alpha-1} \\ s_2 \otimes t_{2\alpha-1} \\ s_1 \otimes t_{2\alpha} \\ s_2 \otimes t_{2\alpha} \end{pmatrix},$$

$$\frac{1}{2\pi}\nabla^2\begin{pmatrix} s_1 \otimes t_{2n+3} \\ s_2 \otimes t_{2n+3} \end{pmatrix} = \begin{pmatrix} 0 & -x \\ x & 0 \end{pmatrix}\begin{pmatrix} s_1 \otimes t_{2n+3} \\ s_2 \otimes t_{2n+3} \end{pmatrix}.$$

Hence the Euler class of Grassmann manifold $G(2, 2n+3)$ is

$$e(TG(2, 2n+3)) = e(E \otimes F) = x\prod_{\alpha=2}^{n+1}(x^2 - y_\alpha^2) = (n+1)e^{2n+1}(E).$$

The odd dimensional homology groups of Grassmann manifold $G(2, 2n+3)$ are also trivial, the even dimensional homology groups are one dimensional. The Euler-Poincaré number is $\chi(G(2, 2n+3)) = 2n+2$.

Similar to the case of $G(2, 2n+2)$, we have

Theorem 5.4 (1) The Pontrjagin classes of $F(2, 2n+3)$ and $TG(2, 2n+3)$ can all be represented by the Euler class $e(E(2, 2n+3))$;

(2) $e(TG(2, 2n+3)) = (n+1)e^{2n+1}(E(2, 2n+3))$;

(3) $\int_{G(2,k+2)} e^k(E(2, 2n+3)) = 2, \quad k = 1, \cdots, 2n+1$;

(4) $\int_{\mathbf{C}P^k} e^k(E(2, 2n+3)) = (-1)^k, \quad k = 1, \cdots, n+1$.

As is well-known, the Chern, Pontrjagin and Euler classes are all integral cocycles. Let D: $H^k(G(2, N), \mathbf{Z}) \to H_{2N-4-k}(G(2, N), \mathbf{Z})$ be the Poincaré duality. The following theorem gives the structure of the integral homology and coholomogy of $G(2, N)$.

Theorem 5.5 (1) When $2k+2 < N$, $[\mathbf{C}P^k]$ and $e^k(E(2, N))$ are the generators of $H_{2k}(G(2, N), \mathbf{Z})$ and $H^{2k}(G(2, N), \mathbf{Z})$ respectively;

(2) When $2k+2>N$, $[G(2,k+2)]$ and $\frac{1}{2}e^k(E(2,N))$ are the generators of $H_{2k}(G(2,N),\mathbf{Z})$ and $H^{2k}(G(2,N),\mathbf{Z})$ respectively;

(3) When $2k+2<N$, $D(e^k(E(2,N)))=[G(2,N-k)]$; when $2k+2>N$, $D(\frac{1}{2}e^k(E(2,N)))=(-1)^{n-k}[\mathbf{C}P^{n-k-2}]$;

(4) $[\mathbf{C}P^n], [\overline{\mathbf{C}P}^n]$ and $\frac{1}{2}\{(-1)^n e^n(E(2,2n+2))\pm e(F(2,2n+2))\}$ are generators of $H_{2n}(G(2,2n+2),\mathbf{Z})$ and $H^{2n}(G(2,2n+2),\mathbf{Z})$ respectively. Furthermore,

$$D(\frac{1}{2}[(-1)^n e^n(E(2,2n+2))+e(F(2,2n+2))])=[\mathbf{C}P^n],$$

$$D(\frac{1}{2}[(-1)^n e^n(E(2,2n+2))-e(F(2,2n+2))])=[\overline{\mathbf{C}P}^n].$$

Proof As is well-known, the Euler classes and Pontrjagin classes are harmonic forms and are integral cocyles, their product are also harmonic forms, see [7], [13]. When $2k+2<N$, from $\int_{\mathbf{C}P^k} e^k(E(2,N))=(-1)^k$ we know $\mathbf{C}P^k\in H_{2k}(G(2,N),\mathbf{Z})$ and $e^k(E(2,N))\in H^{2k}(G(2,N),\mathbf{Z})$ are generators respectively.

By simple computation, we have

$$e^k(E(2,N))=\frac{k!}{(2\pi)^k}\sum_{\alpha_1<\cdots<\alpha_k}\omega_1^{\alpha_1}\omega_2^{\alpha_1}\cdots\omega_1^{\alpha_k}\omega_2^{\alpha_k},$$

$$a=(e^k(E(2,N)),e^k(E(2,N)))=\frac{(k!)^2}{(2\pi)^{2k}}C^k_{N-2}V(G(2,N)),$$

$$\begin{aligned}\frac{1}{a}*e^k(E(2,N)) &= \frac{k!}{a(2\pi)^k}\sum_{\beta_1<\cdots<\beta_{N-k-2}}\omega_1^{\beta_1}\omega_2^{\beta_1}\cdots\omega_1^{\beta_{N-k-2}}\omega_2^{\beta_{N-k-2}}\\ &= \frac{(N-k-2)!}{2(2\pi)^{N-k-2}}\sum_{\beta_1<\cdots<\beta_{N-k-2}}\omega_1^{\beta_1}\omega_2^{\beta_1}\cdots\omega_1^{\beta_{N-k-2}}\omega_2^{\beta_{N-k-2}}\\ &= \frac{1}{2}e^{N-k-2}(E(2,N)).\end{aligned}$$

By Theorem 3.1, $\frac{1}{2}e^{N-k-2}(E(2,N))$ is a generator of $H^{2N-2k-4}(G(2,N),\mathbf{Z})$. By $\int_{G(2,N-k)}\frac{1}{2}e^{N-k-2}(E(2,N))=1$ we know that $G(2,N-k)\in H_{2N-2k-4}(G(2,N),\mathbf{Z})$ is a generator and $D(e^k(E(2,N)))=G(2,N-k)$. These proves (1), (2), (3) of the Theorem.

Let $[S_1],[S_2]$ be generators of $H_{2n}(G(2,2n+2),\mathbf{Z})$ and harmonic forms ξ_1,ξ_2 be generators of $H_{2n}(G(2,2n+2),\mathbf{Z})$, they satisfy $\int_{S_i}\xi_j=\delta_{ij}$. There are integers a_{ij},n_{ij} such that

$$\begin{pmatrix} e^n(E(2,2n+2))\\ e(F(2,2n+2))\end{pmatrix}=\begin{pmatrix} n_{11} & n_{12}\\ n_{21} & n_{22}\end{pmatrix}\begin{pmatrix}\xi_1\\ \xi_2\end{pmatrix},$$

$$(\mathbf{C}P^n, \overline{\mathbf{C}P}^n) = (S_1, S_2)\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

Then

$$\begin{pmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{pmatrix}\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} (-1)^n & (-1)^n \\ 1 & -1 \end{pmatrix},$$

we have $\det(a_{ij}) = \pm 1$ or $\det(n_{ij}) = \pm 1$.

Assuming $\det(n_{ij}) = \pm 1$, then $e^n(E(2, 2n+2)), e(F(2, 2n+2))$ are the generators of $H^{2n}(G(2, 2n+2), \mathbf{Z})$, we can assume $\xi_1 = e^n(E(2, 2n+2)), \xi_2 = e(F(2, 2n+2))$. It is easy to see

$$(e^n(E(2, 2n+2)), e(F(2, 2n+2))) = 0,$$

$$(e^n(E(2, 2n+2)), e^n(E(2, 2n+2))) = (e(F(2, 2n+2)), e(F(2, 2n+2))) = 2.$$

By Theorem 3.1, $\frac{1}{2}e^n(E(2, 2n+2))$, $\frac{1}{2}e(F(2, 2n+2))$ are also the generators of $H^{2n}(G(2, 2n+2), \mathbf{Z})$. This contradict to $e^n(E(2, 2n+2)), e(F(2, 2n+2))$ be the generators.

Then we must have $\det(n_{ij}) = \pm 2$ and $\det(a_{ij}) = \pm 1$. These shows $\mathbf{C}P^n, \overline{\mathbf{C}P}^n$ are generators of $H_{2n}(G(2, 2n+2), \mathbf{Z})$, $\frac{1}{2}\{(-1)^n e^n(E(2, 2n+2)) + e(F(2, 2n+2))\}$ and $\frac{1}{2}\{(-1)^n e^n(E(2, 2n+2)) - e(F(2, 2n+2))\}$ are generators of $H^{2n}(G(2, 2n+2), \mathbf{Z})$. The Poincaré duals of these generators are easy to compute. $\square$

Finally we give some applications to conclude this section.

Let f: $M \to \mathbf{R}^N$ be an immersion of an oriented compact surface M, g: $M \to G(2, N)$ the induced Gauss map, $g(p) = T_pM$. Then $e(M) = g^*(e(E(2, N))$ is the Euler class of M. Let $[M] \in H_2(M)$ be the fundamental class of M. When $N \neq 4$, we have

$$g_*[M] = \frac{1}{2}\chi(M)[G(2, 3)] = \chi(M)[-\mathbf{C}P^1] \in H_2(G(2, N)).$$

In [15], [16], we have shown there is a fibre bundle τ: $G(2, 8) = G(6, 8) \to S^6$ with fibres $\mathbf{C}P^3$, where $S^6 = \{v \in S^7 \mid v \perp \bar{e}_1 = (1, 0, \cdots, 0)\}$, $\tau^{-1}(\bar{e}_2) = \{v \wedge Jv \mid v \in S^7\}$, $\bar{e}_2 = (0, 1, 0, \cdots, 0)$. On the other hand, the map $f(v) = \bar{e}_1 \wedge v$ gives a section of τ. Let $\mathrm{d}V$ be the volume form on S^6 such that $\int_{S^6} \mathrm{d}V = 1$. It is easy to see

$$[\tau^*\mathrm{d}V] = \frac{1}{2}e^3(E(2, 8)) + \frac{1}{2}e(F(2, 8)).$$

Let φ: $M \to \mathbf{R}^8$ be an immersion of an oriented compact 6 dimensional manifold, g: $M \to G(6, 8) = G(2, 8)$ be Gauss map. Then $e(M) = g^*e(F(2, 8))$ is the Euler class of tangent bundle of M, $e(T^{\perp}M) = g^*e(E(2, 8))$ is the Euler class of normal bundle of M.

$$\int_M (\tau \circ g)^*\mathrm{d}V = \int_M g^*[\frac{1}{2}e^3(E(2, 8)) + \frac{1}{2}e(F(2, 8))] = \frac{1}{2}\int_M e^3(T^{\perp}M) + \frac{1}{2}\chi(M)$$

is the degree of the map $\tau \circ g$: $M \to S^6$. If φ is an imbedding, $e(T^{\perp}M) = 0$, see Milnor, Stasheff [12], p.120.

Let J, $\bar{J}$ be two complex structures on $\mathbf{R}^4$, with orthonormal basis e_1, e_2, e_3, e_4,

$$Je_1 = e_2,\ Je_2 = -e_1,\ Je_3 = e_4,\ Je_4 = -e_3;$$

$$\bar{J}e_1 = -e_2,\ \bar{J}e_2 = e_1,\ \bar{J}e_3 = e_4,\ \bar{J}e_4 = -e_3.$$

For any unit vector $v = \sum\ v_i e_i$, we have

$$vJv + *vJv = e_1e_2 + e_3e_4,$$

$$\begin{aligned} vJv - *vJv &= (v_1^2 + v_2^2 - v_3^2 - v_4^2)(e_1e_2 - e_3e_4) \\ &\quad +2(v_1v_3 + v_2v_4)(e_1e_4 - e_2e_3) + 2(v_2v_3 - v_1v_4)(e_1e_3 + e_2e_4); \end{aligned}$$

$$v\bar{J}v - *v\bar{J}v = -e_1e_2 + e_3e_4,$$

$$\begin{aligned} v\bar{J}v + *v\bar{J}v &= (-v_1^2 - v_2^2 + v_3^2 + v_4^2)(e_1e_2 + e_3e_4) \\ &\quad +2(v_1v_3 - v_2v_4)(e_1e_4 + e_2e_3) - 2(v_1v_4 + v_2v_3)(e_1e_3 - e_2e_4). \end{aligned}$$

These shows $\mathbf{C}P^1$, $\overline{\mathbf{C}P}^1$ are two spheres in $G(2,4) \approx S^2(\frac{\sqrt{2}}{2}) \times S^2(\frac{\sqrt{2}}{2})$ where the decomposition is given by Hodge star operator. Let f: $M \to \mathbf{R}^4$ be an imbedding of an oriented surface, g: $M \to G(2,4)$ the Gauss map. Then we have

$$g_*[M] = \frac{1}{2}\chi(M)[G(2,3)] = -\frac{1}{2}\chi(M)[\mathbf{C}P^1] - \frac{1}{2}\chi(M)[\overline{\mathbf{C}P}^1].$$

See the work of Chern and Spanier [1].

§6. The case of $G(3,6)$

The Poincaré polynomial of $G(3,6)$ is $p_t(G(3,6)) = 1 + t^4 + t^5 + t^9$. To study the real homology and coholomogy of Grassmann manifold $G(3,6)$ we need only consider the dimension 4 and 5.

Let i: $G(2,4) \to G(3,6)$ be inclusion defined naturally. It is easy to see that $i^*p_1(E(3,6)) = p_1(E(2,4)) = e^2(E(2,4))$, then

$$\int_{G(2,4)} p_1(E(3,6)) = 2.$$

As §2, let $SLAG = \{G(\bar{e}_1\bar{e}_3\bar{e}_5) \mid G \in SU(3) \subset SO(6)\}$ be a subspace of $G(3,6)$, $e_i = G(\bar{e}_i), e_{i+1} = G(\bar{e}_{i+1}) = Je_i$ be $SU(3)$-frame fields, $i = 1,3,5$. Restricting the coframes $\omega_A^B = \langle \mathrm{d}e_A, e_B\rangle$ on $SLAG$ we have

$$\omega_i^j = \omega_{i+1}^{j+1},\ \omega_i^{j+1} = -\omega_{i+1}^j,\ i,j = 1,3,5 \ \text{ and } \ \omega_1^2 + \omega_3^4 + \omega_5^6 = 0.$$

By the proof of Proposition 2.4, we have $\mathrm{d}V_{SLAG} = 2^{\frac{3}{2}}\sqrt{3}\,\omega_1^4\omega_1^6\omega_3^6\omega_1^2\omega_3^4$ and $V(SLAG) = \sqrt{\frac{3}{2}}\,\pi^3$. Let $G(3,6)$ be generated by $e_1e_3e_5$ locally, the first Pontrjagin class of canonical vector bundle $E(3,6)$ is

$$p_1(E(3,6)) = \frac{1}{4\pi^2}[(\Omega_{13})^2 + (\Omega_{15})^2 + (\Omega_{35})^2],$$

where $\Omega_{ij} = -\sum\limits_{\alpha} \ \omega_i^\alpha \wedge \omega_j^\alpha,\ \alpha = 2,4,6$. By computation we have

$$*p_1(E(3,6))|_{SLAG} = \frac{\sqrt{6}}{4\pi^2}\mathrm{d}V_{SLAG},$$

$$a = (p_1(E(3,6)), p_1(E(3,6))) = \frac{3\cdot 4\cdot 3}{(2\pi)^4}V(G(3,6)) = \frac{3}{2}\pi.$$

From $\int_{G(2,4)} \ p_1(E(3,6)) = 2$, we know that $p_1(E(3,6))$ or $\frac{1}{2}p_1(E(3,6))$ is a generator of $H^4(G(3,6),\mathbf{Z})$. If $p_1(E(3,6))$ is a generator, by Theorem 3.1, $\frac{1}{a} * p_1(E(3,6))$ is a generator of $H^5(G(3,6),\mathbf{Z})$, but

$$\int_{SLAG} \ \frac{1}{a} * p_1(E(3,6)) = \int_{SLAG} \ \frac{1}{\sqrt{6}\pi^3}\mathrm{d}V_{SLAG} = \frac{1}{2}.$$

Then $\frac{1}{2}p_1(E(3,6))$ is a generator of $H^4(G(3,6),\mathbf{Z})$ and $\int_{SLAG} \ \frac{4}{a} * \frac{1}{2}p_1(E(3,6)) = 1$.

We have proved following theorem

Theorem 6.1 (1) $\frac{1}{2}p_1(E(3,6)) \in H^4(G(3,6),\mathbf{Z})$ is a generator and its Poincaré dual $[SLAG]$ is a generator of $H_5(G(3,6),\mathbf{Z})$;

(2) $\frac{4}{3\pi} * p_1(E(3,6)) \in H^5(G(3,6),\mathbf{Z})$ is a generator and its Poincaré dual $[G(2,4)]$ is a generator of $H_4(G(3,6),\mathbf{Z})$.

Let $\bar{e}_1,\cdots,\bar{e}_6$ be a fixed orthonormal basis of $\mathbf{R}^6$, $G \in SO(3)$ acts on the subspace generated by $\bar{e}_4,\bar{e}_5,\bar{e}_6$, denote $e_4 = G(\bar{e}_4), e_5 = G(\bar{e}_5), e_6 = G(\bar{e}_6)$. As [7], let $PONT$ be the set of elements

$$(\cos t\bar{e}_1 + \sin te_4)(\cos t\bar{e}_2 + \sin te_5)(\cos t\bar{e}_3 + \sin te_6), \quad t \in [0, \frac{\pi}{2}].$$

$PONT$ is a calibrated submanifold (except two points correspond to $t = 0, \frac{\pi}{2}$) of the first Pontrjagin form $p_1(G(3,6))$. By moving frame we can show

$$\int_{PONT} \ p_1(G(3,6)) = 2 \ \text{ and } \ V(PONT) = \sqrt{\frac{2}{3}}\,V(G(2,4)) = \frac{4\sqrt{6}}{3}\pi^2.$$

Then 4-cycle $PONT$ is homologous to the 4-cycle $G(2,4)$ inside $G(3,6)$.

§7. The case of $G(3,7)$

The Poincaré polynomial of $G(3,7)$ is $p_t(G(3,7)) = 1 + 2t^4 + 2t^8 + t^{12}$.

Let $\bar{e}_1, \bar{e}_2, \cdots, \bar{e}_8$ be a fixed orthonormal basis of $\mathbf{R}^8$ and $\mathbf{R}^7$ be a subspace generated by $\bar{e}_2, \cdots, \bar{e}_8$. The oriented Grassmann manifold $G(3,7)$ is the set of subspaces of $\mathbf{R}^7$.

Let $E = E(3,7)$ and $F = F(3,7)$ be canonical vector bundles on Grassmann manifold $G(3,7)$. As §4, we can show

$$p_1(F) = -p_1(E),\ p_2(F) = e^2(F) = p_1^2(E),\ e(E \otimes F) = e(TG(3,7)) = 3e^3(F).$$

By $\int_{G(3,7)}\ e(TG(3,7)) = \chi(G(3,7)) = 6$ we have

$$\int_{G(3,7)}\ e^3(F) = 2.$$

By inclusion $G(2,6) \subset G(3,7)$, $\mathbf{C}P^2$ and $\overline{\mathbf{C}P}^2$ can be imbedded in $G(3,7)$.

Lemma 7.1 $\int_{\mathbf{C}P^2}\ p_1(E) = \int_{\overline{\mathbf{C}P}^2}\ p_1(E) = 1,\quad \int_{\mathbf{C}P^2}\ e(F) = -\int_{\overline{\mathbf{C}P}^2}\ e(F) = 1.$

Proof By $p_1(E(3,7))|_{\mathbf{C}P^2} = p_1(E(2,6))|_{\mathbf{C}P^2} = e^2(E(2,6))|_{\mathbf{C}P^2}$ and the results of §5, we have $\int_{\mathbf{C}P^2}\ p_1(E) = 1$. The other equalities can be proved similarly. □

Then

$$\int_{\mathbf{C}P^2}\ \frac{1}{2}(p_1(E) + e(F)) = \int_{\overline{\mathbf{C}P}^2}\ \frac{1}{2}(p_1(E) - e(F)) = 1,$$
$$\int_{\mathbf{C}P^2}\ \frac{1}{2}(p_1(E) - e(F)) = \int_{\overline{\mathbf{C}P}^2}\ \frac{1}{2}(p_1(E) + e(F)) = 0,$$

hence $\mathbf{C}P^2, \overline{\mathbf{C}P}^2 \in H_4(G(3,7))$ and $\frac{1}{2}(p_1(E)+e(F)),\ \frac{1}{2}(p_1(E)-e(F)) \in H^4(G(3,7))$ are generators.

Let $e_2, e_3, e_4, \cdots, e_8$ be oriented orthonormal frame fields on $\mathbf{R}^7$, $G(3,7)$ be generated by $e_2 \wedge e_3 \wedge e_4$ locally. Euler class of F and first Pontrjagin class of E can be represented by

$$\begin{aligned} e(F) &= \frac{1}{2(4\pi)^2} \sum\ \varepsilon(\alpha_1\alpha_2\alpha_4\alpha_4)\Omega_{\alpha_1\alpha_2} \wedge \Omega_{\alpha_3\alpha_4} \\ &= \frac{1}{4\pi^2} \sum_{i,j=2}^{4}\ (\omega_5^i\omega_6^i\omega_7^j\omega_8^j - \omega_5^i\omega_7^i\omega_6^j\omega_8^j + \omega_5^i\omega_8^i\omega_6^j\omega_7^j), \end{aligned}$$

$$p_1(E) = \frac{1}{4\pi^2}[(\Omega_{23})^2 + (\Omega_{24})^2 + (\Omega_{34})^2].$$

Lemma 7.2 (1) $*p_1(E) = \frac{4}{5}\pi^2 p_1(E)e(F),\ *e(F) = \frac{1}{2}\pi^2 e^2(F)$;

(2) $(p_1(E), e(F)) = 0,\ a = (p_1(E), p_1(E)) = \frac{8}{5}\pi^2,\ b = (e(F), e(F)) = \pi^2$.

Proof $*p_1(E),\ p_1(E)e(F)$ and $*e(F),\ e^2(F)$ are the harmonic forms on $G(3,7)$. $*p_1(E) = \frac{4}{5}\pi^2 p_1(E)e(F)$ follows from the equalities such as

$$\begin{aligned} *\omega_2^5\omega_3^5\omega_2^6\omega_3^6 &= \omega_2^7\omega_3^7\omega_2^8\omega_3^8\omega_5^4\omega_6^4\omega_7^4\omega_8^4 \\ &= \omega_3^7\omega_4^7\omega_3^8\omega_4^8\omega_5^4\omega_6^4\omega_7^2\omega_8^2 \\ &= \omega_2^7\omega_4^7\omega_2^8\omega_4^8\omega_5^4\omega_6^4\omega_7^3\omega_8^3. \end{aligned}$$

The proof of (2) is a direct computation. □

To study $G(3,7), G(3,8)$ and $G(4,8)$, we shall use Clifford algebras.

Let $C\ell_8$ be the Clifford algebra associated to the Euclidean space $\mathbf{R}^8$. Let $\bar{e}_1, \bar{e}_2, \cdots, \bar{e}_8$ be a fixed orthonormal basis of $\mathbf{R}^8$, the Clifford product be determined by the relations: $\bar{e}_B \cdot \bar{e}_C + \bar{e}_C \cdot \bar{e}_B = -2\delta_{BC},\quad B, C = 1, 2, \cdots, 8$. Define the subspace $V = V^+ \oplus V^-$ of $C\ell_8$ by $V^+ = C\ell_8^{even} \cdot A, V^- = C\ell_8^{odd} \cdot A$, where

$$A = \frac{1}{16}\mathrm{Re}\,[(\bar{e}_1 + \sqrt{-1}\bar{e}_2)\cdots(\bar{e}_7 + \sqrt{-1}\bar{e}_8)(1 + \bar{e}_1\bar{e}_3\bar{e}_5\bar{e}_7)].$$

The space $V = V^+ \oplus V^-$ is an irreducible module over $C\ell_8$. The spaces V^+ and V^- are generated by $\bar{e}_1\bar{e}_B A$ and $\bar{e}_B A$ respectively, $B = 1, \cdots, 8$, see [14], [15].

Let $Spin_7 = \{G \in SO(8) \mid G(A) = A\}$ be the isotropy group of $SO(8)$ acting on A. The group $Spin_7$ acts on $G(2,8), G(3,8)$ and S^7 transitively. $G_2 = \{G \in Spin_7 \mid G(\bar{e}_1) = \bar{e}_1\}$ is a subgroup of $Spin_7$.

The Grassmann manifold $G(k,8)$ can be viewed as subset of Clifford algebra $C\ell_8$ naturally. Then for any $\pi \in G(k,8)$, there is $v \in \mathbf{R}^8$ such that $\pi A = \bar{e}_1 v A$ or $\pi A = vA$ according to the number k being even or odd, $|v| = 1$. Thus we have maps $G(k,8) \to S^7,\ \pi \mapsto v$. Since $Spin_7$ acts on $G(3,8)$ transitively, from $\bar{e}_2\bar{e}_3\bar{e}_4 A = \bar{e}_1 A$ we have $G(\bar{e}_2\bar{e}_3\bar{e}_4)A = G(\bar{e}_1)A$ for any $G \in Spin_7$. This shows the map τ: $G(3,8) \to S^7,\ \tau(\pi) = v$, is a fibre bundle and $v \perp \pi$, see [15], [16]. Let

$$ASSOC = \tau^{-1}(\bar{e}_1) = \{\pi \in G(3,8) \mid \tau(\pi) = \bar{e}_1\}$$

be the fibre over $\bar{e}_1$. The group G_2 acts on $ASSOC$ transitively, we have $ASSOC = \{G(\bar{e}_2\bar{e}_3\bar{e}_4) \mid G \in G_2\}$. We can show the isotropy group $\{G(\bar{e}_2\bar{e}_3\bar{e}_4) = \bar{e}_2\bar{e}_3\bar{e}_4 \mid G \in G_2\}$ is isomorphic to the group $SO(4)$, then $ASSOC \approx G_2/SO(4)$.

Change the orientation of $\mathbf{R}^7$, let $\tilde{A} = \frac{1}{16}\mathrm{Re}\,[(\bar{e}_1 - \sqrt{-1}\bar{e}_2)(\bar{e}_3 + \sqrt{-1}\bar{e}_4)\cdots(\bar{e}_7 + \sqrt{-1}\bar{e}_8)(1 + \bar{e}_1\bar{e}_3\bar{e}_5\bar{e}_7)]$. Define submanifold $\widetilde{ASSOC} = \{\pi \in G(3,8) \mid \pi\tilde{A} = \bar{e}_1\}$ which diffeomorphic to $ASSOC$.

Lemma 7.3 $V(ASSOC) = \frac{6}{5}\pi^4$.

Proof Let $\tilde{e}_1, \tilde{e}_2, \cdots, \tilde{e}_8$ be $Spin_7$ frame fields on $\mathbf{R}^8$, the 1-forms $\omega_B^C = \langle \mathrm{d}\tilde{e}_B, \tilde{e}_C \rangle$ satisfy (for proof, see [16])

$$\begin{array}{ll} \omega_1^2 + \omega_3^4 + \omega_5^6 + \omega_7^8 = 0, & \omega_1^3 - \omega_2^4 + \omega_6^8 - \omega_5^7 = 0, \\ \omega_1^4 + \omega_2^3 + \omega_5^8 + \omega_6^7 = 0, & \omega_1^5 - \omega_2^6 + \omega_3^7 - \omega_4^8 = 0, \\ \omega_1^6 + \omega_2^5 - \omega_3^8 - \omega_4^7 = 0, & \omega_1^7 - \omega_2^8 - \omega_3^5 + \omega_4^6 = 0, \\ \omega_1^8 + \omega_2^7 + \omega_3^6 + \omega_4^5 = 0. & \end{array}$$

Since $Spin_7$ acts on $G(3,8)$ transitively, $G(3,8)$ is locally generated by $\tilde{e}_2\tilde{e}_3\tilde{e}_4$. The volume element of $G(3,8)$ is $\mathrm{d}V_{G(3,8)} = \omega_2^1\omega_3^1\omega_4^1\omega_2^5\omega_3^5\omega_4^5 \cdots \omega_2^8\omega_3^8\omega_4^8$.

Note that A can be represented by $Spin_7$ frames, that is

$$A = \frac{1}{16}\mathrm{Re}\,[(\tilde{e}_1 + \sqrt{-1}\tilde{e}_2)\cdots(\tilde{e}_7 + \sqrt{-1}\tilde{e}_8)(1 + \tilde{e}_1\tilde{e}_3\tilde{e}_5\tilde{e}_7)].$$

Let $\tilde{e}_1 = \bar{e}_1$ be a fixed vector, $\bar{e}_1, \tilde{e}_2, \cdots, \tilde{e}_8$ be G_2 frame fields on $\mathbf{R}^8$, $ASSOC$ is locally generated by $\tilde{e}_2\tilde{e}_3\tilde{e}_4$ and

$$\begin{aligned} \mathrm{d}(\tilde{e}_2\tilde{e}_3\tilde{e}_4) &= \sum_{i=2}^{3}\sum_{\alpha=5}^{8} \omega_i^\alpha E_{i\alpha} \\ &= \omega_2^5(E_{25} + E_{47}) + \omega_2^6(E_{26} - E_{48}) + \omega_2^7(E_{27} - E_{45}) + \omega_2^8(E_{28} + E_{46}) \\ &\quad + \omega_3^5(E_{35} + E_{46}) + \omega_3^6(E_{36} - E_{45}) + \omega_3^7(E_{37} + E_{48}) + \omega_3^8(E_{38} - E_{47}). \end{aligned}$$

The metric on $ASSOC$ is

$$\begin{aligned} \mathrm{d}s^2 &= 2(\omega_2^5)^2 + 2(\omega_3^8)^2 - 2\omega_2^5\omega_3^8 + 2(\omega_2^6)^2 + 2(\omega_3^7)^2 - 2\omega_2^6\omega_3^7 \\ &\quad + 2(\omega_2^7)^2 + 2(\omega_3^6)^2 + 2\omega_2^7\omega_3^6 + 2(\omega_2^8)^2 + 2(\omega_3^5)^2 + 2\omega_2^8\omega_3^5, \end{aligned}$$

with the volume form

$$\mathrm{d}V_{ASSOC} = 9\ \omega_2^5\omega_3^5\ \cdots\ \omega_2^8\omega_3^8.$$

The normal space of $ASSOC$ in $G(3,8)$ at $\tilde{e}_2\tilde{e}_3\tilde{e}_4$ is generated by

$$E_{21}, E_{31}, E_{41}, E_{25} - E_{47} - E_{38}, E_{26} + E_{48} - E_{37}, E_{27} + E_{45} + E_{36}, E_{28} - E_{46} + E_{35}.$$

The sphere S^7 is generated by $\tilde{e}_1$, $\mathrm{d}V_{S^7} = \omega_2^1\omega_3^1 \cdots \omega_1^8$ is the volume form. From

$$(E_{27} + E_{45} + E_{36})A = -3\tilde{e}_8 A,\ E_{21}A = -\tilde{e}_2 A,\ \cdots,$$

we can compute the tangent map of τ: $G(3,8) \to S^7$,

$$\tau_*(E_{27} + E_{45} + E_{36}) = -3\tilde{e}_8,\ \tau_*(E_{21}) = -\tilde{e}_2,\ \cdots.$$

Then we can compute the cotangent map τ^* and we have

$$\tau^*\mathrm{d}V_{S^7} = \omega_2^1\omega_3^1\omega_4^1(\omega_2^6 - \omega_3^7 + \omega_4^8)(-\omega_2^5 + \omega_3^8 + \omega_4^7)(\omega_2^8 + \omega_3^5 - \omega_4^6)(\omega_2^7 + \omega_3^6 + \omega_4^5),$$

and

$$\mathrm{d}V_{G(3,8)} = -\frac{1}{9}\tau^*\mathrm{d}V_{S^7} \cdot \mathrm{d}V_{\tau^{-1}(\tilde{e}_1)}.$$

From $V(G(3,8)) = \frac{2}{45}\pi^8,\ V(S^7) = \frac{1}{3}\pi^4$, we have $V(ASSOC) = \frac{6}{5}\pi^4$. $\square$

Lemma 7.4 $\int_{ASSOC}\ p_1^2(E) = \int_{ASSOC}\ p_1(E)e(F) = 1$,

$\int_{\widetilde{ASSOC}}\ p_1^2(E) = -\int_{\widetilde{ASSOC}}\ p_1(E)e(F) = 1$.

Then $ASSOC, \widetilde{ASSOC}$ and $p_1^2(E), p_1(E)e(F)$ are generators of $H_8(G(3,7))$ and $H^8(G(3,7))$ respectively. Furthermore, we have

$$[G(2,6)] = [ASSOC] + [\widetilde{ASSOC}].$$

Proof From

$$*\omega_2^5\omega_3^5\omega_2^6\omega_3^6|_{ASSOC} = \omega_2^7\omega_3^7\omega_2^8\omega_3^8\omega_5^4\omega_6^4\omega_7^4\omega_8^4|_{ASSOC} = \frac{1}{9}\mathrm{d}V_{ASSOC},$$

$$*\omega_2^7\omega_3^7\omega_2^8\omega_3^8|_{ASSOC} = *\omega_2^5\omega_3^5\omega_2^7\omega_3^7|_{ASSOC} = *\omega_2^6\omega_3^6\omega_2^8\omega_3^8|_{ASSOC} = \frac{1}{9}\mathrm{d}V_{ASSOC},$$

$$*\omega_2^6\omega_3^6\omega_2^7\omega_3^7|_{ASSOC} = \omega_2^5\omega_3^5\omega_2^8\omega_3^8\omega_5^4\omega_6^4\omega_7^4\omega_8^4|_{ASSOC} = 0,$$

$$*\omega_2^5\omega_3^5\omega_2^8\omega_3^8|_{ASSOC} = 0,$$

we have $\sum\limits_{\alpha,\beta} *\omega_2^\alpha\omega_3^\alpha\omega_2^\beta\omega_3^\beta|_{ASSOC} = 8 \cdot \frac{1}{9}\mathrm{d}V_{ASSOC}$,

$$*p_1(E)|_{ASSOC} = \frac{1}{4\pi^2} \cdot 3 \cdot 8 \cdot \frac{1}{9}\mathrm{d}V_{ASSOC}.$$

Then by $*p_1(E) = \frac{4}{5}\pi^2 p_1(E)e(F)$, we have

$$\int_{ASSOC} p_1(E)e(F) = \int_{ASSOC} \frac{5}{4\pi^2} * p_1(E) = 1.$$

The proof of $\int_{ASSOC}\ p_1^2(E) = 1$ is similar. $\square$

Change the orientation of $\mathbf{R}^7$, we have Euclidean space $\widetilde{\mathbf{R}}^7$. Let $\widetilde{E}, \widetilde{F} \to G(3,7)$ be canonical vector bundles with respect to $\widetilde{\mathbf{R}}^7$. It is easy to see that $\widetilde{E} = E$, but the orientations of $\widetilde{F}$ and F are different. These shows

$$\int_{\widetilde{ASSOC}} p_1^2(E) = 1, \quad \int_{\widetilde{ASSOC}} p_1(E)e(F) = -1.$$

$$[G(2,6)] = [ASSOC] + [AS\widetilde{SOC}]$$

follows from $\int_{G(2,6)} p_1^2(E) = 2$ and $\int_{G(2,6)} p_1(E)e(F) = 0$.

Theorem 7.5 (1) $\frac{1}{2}(p_1(E) + e(F))$, $\frac{1}{2}(p_1(E) - e(F))$ are two generators of $H^4(G(3,7),\mathbf{Z})$. Their Poincaré duals are $[ASSOC]$ and $[AS\widetilde{SOC}]$ respectively;

(2) $\frac{1}{2}(p_1(E)e(F)+e^2(F))$, $\frac{1}{2}(p_1(E)e(F)-e^2(F)) \in H^8(G(3,7),\mathbf{Z})$ are generators and their Poincaré duals are $[\mathbf{C}P^2]$, $[\overline{\mathbf{C}P}^2]$ respectively;

(3) $\frac{1}{2}(p_1(E)e(F) + e^2(F))$, $\frac{1}{2}(p_1(E)e(F) - e^2(F))$ and $[ASSOC]$, $[AS\widetilde{SOC}]$ are dual basis with respect to the universal coefficients Theorem.

The proof is similar to Theorem 5.5.

§8. The case of $G(3,8)$

The Poincaré polynomial of Grassmann manifold $G(3,8)$ is

$$p_t(G(3,8)) = (1 + t^4 + t^8)(1 + t^7) = 1 + t^4 + t^7 + t^8 + t^{11} + t^{15}.$$

Let $E = E(3,8)$, $F = F(3,8)$ be canonical vector bundles on $G(3,8)$. By $\int_{\mathbf{C}P^2} p_1(E) = 1$, $\int_{ASSOC} p_1^2(E) = 1$ we know that

$$[\mathbf{C}P^2] \in H_4(G(3,8),\mathbf{Z}),\ [ASSOC] \in H_8(G(3,8),\mathbf{Z}),$$

$$p_1(E) \in H^4(G(3,8),\mathbf{Z}),\ p_1^2(E) \in H^8(G(3,8),\mathbf{Z})$$

are all generators. By Theorem 3.1, to understand the structure of the homology and coholomogy groups of dimension $7, 11$, we need compute the Poincaré duals of $p_1(E)$, $p_1^2(E)$.

It is not difficult to compute

$$a = (p_1(E), p_1(E)) = \frac{15}{2\pi^4}V(G(3,8)) = \frac{1}{3}\pi^4.$$

Then $\frac{1}{a} * p_1(E) \in H^{11}(G(3,8),\mathbf{Z})$ is a generator, we look for a submanifold M such that $\int_M \frac{1}{a} * p_1(E) = 1$.

In §7 we define fibre bundle τ: $G(3,8) \to S^7$, $\tau(\pi) = v$ defined by $\pi A = vA$. As $v \perp \pi$, $v\pi \in G(4,8)$ and $v\pi A = -A$. Let $CAY = \{\pi \in G(4,8) \mid \pi A = -A\}$, called Cayley submanifold of $G(4,8)$. Then we have a fibre bundle μ: $G(3,8) \to CAY$, $\pi \mapsto v\pi$, with fibre $S^3 = G(3,4)$. Let $ASSOC = \{\bar{e}_1e_2e_3e_4 \mid e_2e_3e_4A = \bar{e}_1A\}$, where $\bar{e}_1 = (1,0,\cdots,0)$. Then $M = \mu^{-1}(ASSOC)$ is a 11 dimensional submanifold of $G(3,8)$.

Let $\bar{e}_1, e_2, \cdots, e_8$ be G_2 frame fields on $\mathbf{R}^8$, $ASSOC$ be generated by $\bar{e}_1 e_2 e_3 e_4$. Represent the elements of $\mu^{-1}(\bar{e}_1 e_2 e_3 e_4)$ by $\tilde{e}_2\tilde{e}_3\tilde{e}_4$, $\tau(\tilde{e}_2\tilde{e}_3\tilde{e}_4) = \tilde{e}_1$, then

$$\tilde{e}_1\tilde{e}_2\tilde{e}_3\tilde{e}_4 = \bar{e}_1 e_2 e_3 e_4.$$

Let $\mathrm{d}(\tilde{e}_2\tilde{e}_3\tilde{e}_4) = \sum\limits_{i=2}^{4} \tilde{\omega}_i^1 \tilde{E}_{i1} + \sum\limits_{i=2}^{4}\sum\limits_{\alpha=5}^{8} \tilde{\omega}_i^\alpha \tilde{E}_{i\alpha}$, $\mathrm{d}s_M^2 = \sum\limits_{i} (\tilde{\omega}_i^1)^2 + \sum\limits_{i,\alpha} (\tilde{\omega}_i^\alpha)^2$ be the metric on M.

Let $\tilde{e}_1 = \lambda_1\bar{e}_1 + \sum\limits_{i=2}^{4} \lambda_i e_i$, $\mathrm{d}\tilde{e}_1 = \sum\limits_{i=2}^{4} \tilde{\omega}_1^i \tilde{e}_i + \sum\limits_{\alpha=5}^{8} \tilde{\omega}_1^\alpha e_\alpha$ and $\mathrm{d}(e_2e_3e_4) = \sum\limits_{i,\alpha} \omega_i^\alpha E_{i\alpha}$. It is easy to see $\tilde{\omega}_1^\alpha = \sum \lambda_i \omega_i^\alpha$. $\sum\limits_{i=2}^{4} (\tilde{\omega}_1^i)^2$ is the metric on the fibres of μ: $M \to ASSOC$ and $\langle \mathrm{d}(e_2e_3e_4), \mathrm{d}(e_2e_3e_4)\rangle = \sum (\omega_i^\alpha)^2$ is metric on $ASSOC$. From $\langle \bar{e}_1 E_{i\alpha}, e_\beta\tilde{e}_2\tilde{e}_3\tilde{e}_4\rangle$ $= \lambda_i\delta_{\alpha\beta}$ we have

$$\langle \bar{e}_1\mathrm{d}(e_2e_3e_4), (\mathrm{d}\tilde{e}_1)\tilde{e}_2\tilde{e}_3\tilde{e}_4\rangle = \langle \sum \omega_i^\alpha \bar{e}_1 E_{i\alpha}, \sum \lambda_j \omega_j^\beta e_\beta \tilde{e}_2\tilde{e}_3\tilde{e}_4\rangle = \sum_\alpha (\sum_i \lambda_i\omega_i^\alpha)^2.$$

Hence

$$\begin{aligned}
\sum (\tilde{\omega}_i^\alpha)^2 &= \langle \tilde{e}_1\mathrm{d}(\tilde{e}_2\tilde{e}_3\tilde{e}_4), \tilde{e}_1\mathrm{d}(\tilde{e}_2\tilde{e}_3\tilde{e}_4)\rangle \\
&= \langle \bar{e}_1\mathrm{d}(e_2e_3e_4) - (\mathrm{d}\tilde{e}_1)\tilde{e}_2\tilde{e}_3\tilde{e}_4, \bar{e}_1\mathrm{d}(e_2e_3e_4) - (\mathrm{d}\tilde{e}_1)\tilde{e}_2\tilde{e}_3\tilde{e}_4\rangle \\
&= \sum_{i,\alpha} (\omega_i^\alpha)^2 + \sum (\tilde{\omega}_1^\alpha)^2 - 2\sum_\alpha (\sum_i \lambda_i\omega_i^\alpha)^2 \\
&= \sum_{i,\alpha} (\omega_i^\alpha)^2 - \sum_\alpha (\sum_i \lambda_i\omega_i^\alpha)^2.
\end{aligned}$$

Then the metric on M can be represented by

$$\mathrm{d}s_M^2 = \sum_{i,\alpha} (\omega_i^\alpha)^2 - \sum_\alpha (\sum_i \lambda_i\omega_i^\alpha)^2 + \sum_{i=2}^{4} (\tilde{\omega}_1^i)^2.$$

For fixed $\tilde{e}_2\tilde{e}_3\tilde{e}_4$ of M, we can choice G_2 frame fields $\bar{e}_1, e_2, \cdots, e_8$ such that $\tilde{e}_1 = \lambda_1\bar{e}_1 + \lambda_2 e_2$. By

$$\omega_4^5 = -\omega_2^7 - \omega_3^6,\ \omega_4^6 = \omega_2^8 + \omega_3^5,\ \omega_4^7 = \omega_2^5 - \omega_3^8,\ \omega_4^8 = -\omega_2^6 + \omega_3^7,$$

we have

$$\mathrm{d}V_M = \frac{1}{9}(1+2\lambda_1^2)^2\tilde{\omega}_1^2\tilde{\omega}_1^3\tilde{\omega}_1^4\mu^*\mathrm{d}V_{ASSOC}.$$

We can show that $\int_{S^3} (1+2\lambda_1^2)^2\mathrm{d}V_{S^3} = 5\pi^2$, then $V(M) = \frac{2}{3}\pi^6$.

Lemma 8.1 $\int_M \frac{3}{\pi^4} * p_1(E) = 1$. Then $[M]$ and $\frac{3}{\pi^4} * p_1(E)$ are dual generators of $H_8(G(3,8),\mathbf{Z})$, $H^8(G(3,8),\mathbf{Z})$ respectively.

Proof By Theorem 3.1, the integration $\int_M \frac{1}{a} * p_1(E)$ is an integer. On the other hand $2\pi^2 p_1(E)$ is a calibration on $G(3,8)$ with comass $\frac{4}{3}$, see [7], then we have

$$\Big|\int_M \frac{1}{a} * p_1(E)\Big| \le \frac{2}{3a\pi^2}V(M) = \frac{4}{3}.$$

We need to show that $\int_M \frac{1}{a} * p_1(E) \neq 0$.

Let i: $CAY \to G(4,8)$ be the inclusion, μ: $G(3,8) \to CAY$ be sphere bundle associated to the induced vector bundle $i^*E(4,8) \to CAY$, $e(i^*E(4,8)) = i^*e(E(4,8)) \in H^4(CAY, \mathbf{Z})$ be the Euler class. The induced bundle $(i\circ\tau)^*E(4,8) \to G(3,8)$ has a nonzero section, $(i\circ\tau)^*e(E(4,8)) = 0$. By Gysin sequence for the sphere bundle $G(3,8) \to CAY$, we can show $\tau_*(\frac{1}{a} * p_1(E)), (i^*e(E(4,8)))^2$ are two generators of $H^8(CAY)$, where τ_*: $H^{11}(G(3,8)) \to H^8(CAY)$ is the integration along the fibre. It is easy to see that $\int_{ASSOC} (i^*e(E(4,8)))^2 = 0$. If we also have $\int_{ASSOC} \tau_*(\frac{1}{a} * p_1(E)) = 0$, then $[ASSOC] = 0$ in $H_8(CAY)$. This contradict to the fact $i_*[ASSOC] \neq 0$. Then with suitable choice orientation on M, we have

$$\int_M \frac{1}{a} * p_1(E) = \int_{ASSOC} \tau_*(\frac{1}{a} * p_1(E)) = 1. \quad \square$$

Finally we study the $H_7(G(3,8))$ and $H^7(G(3,8))$. Let I, J, K be the quaternion structures on $\mathbf{R}^8 = \mathbf{H}^2$ and $Sp(2)$ the symplectic group. As we know $Sp(2)$ can be viewed as a subgroup of $SU(4) \subset SO(8)$, $Sp(2)$ is a subgroup of $Spin_7$. Let f: $S^7 \to G(3,8)$, $f(v) = IvJvKv$. By $I\bar{e}_1 J\bar{e}_1 K\bar{e}_1 A = \bar{e}_2\bar{e}_3\bar{e}_4 A = \bar{e}_1 A$ and $Sp(2)$ acts on S^7 transitively, we have $IvJvKvA = vA$ for any $v \in S^7$, $\tau(f(v)) = v$.

Lemma 8.2 $b = (p_1^2(E), p_1^2(E)) = \frac{5}{2}$.

Proof By computation, we have

$$\begin{aligned} p_1^2(E) &= \frac{1}{2\pi^4}\{\sum_{i<j}\sum_{\alpha<\beta<\gamma<\tau} 3\omega_i^\alpha\omega_j^\alpha\omega_i^\beta\omega_j^\beta\omega_i^\gamma\omega_j^\gamma\omega_i^\tau\omega_j^\tau \\ &\quad + \sum_{j\neq k}\sum_{\alpha<\beta,\gamma<\tau} \omega_i^\alpha\omega_j^\alpha\omega_i^\beta\omega_j^\beta\omega_i^\gamma\omega_k^\gamma\omega_i^\tau\omega_k^\tau\}, \end{aligned}$$

$$i, j, k = 2, 3, 4,\ \ \alpha, \beta, \gamma, \tau = 1, 5, 6, 7, 8.$$

Then

$$b = (p_1^2(E), p_1^2(E)) = \frac{1}{4\pi^8}(9\cdot 3\cdot C_5^4 + 3\cdot C_5^2\cdot C_3^2)V(G(3,8)) = \frac{5}{2}. \quad \square$$

$p_1^2(E)$ is a harmonic form, by Theorem 3.1, $\frac{1}{b} * p_1^2(E) \in H^7(G(3,8), \mathbf{Z})$ is a generator.

Lemma 8.3 $\frac{2}{5}\int_{S^7}\ f^{*}{*}p_1^2(E)=1$. Then $[f(S^7)]$ and $\frac{2}{5}{*}p_1^2(E)$ are dual generators of $H_7(G(3,8),\mathbf{Z}),\ H^7(G(3,8),\mathbf{Z})$ respectively.

Proof Let $e_1, e_2=Ie_1, e_3=Je_1, e_4=Ke_1, e_5, e_6=Ie_5, e_7=Je_5, e_8=Ke_5$ be $Sp(2)$ frame fields on $\mathbf{R}^8$, $f(e_1)=e_2e_3e_4$. The 1 forms $\omega_i^\alpha=\langle \mathrm{d}e_i, e_\alpha\rangle,\ i=1,2,3,4,\ \alpha=5,6,7,8$, satisfy

$$\omega_1^5=\omega_2^6=\omega_3^7=\omega_4^8,\quad \omega_1^6=-\omega_2^5=-\omega_3^8=\omega_4^7,$$

$$\omega_1^7=\omega_2^8=-\omega_3^5=-\omega_4^6,\quad \omega_1^8=-\omega_2^7=\omega_3^6=-\omega_4^5.$$

We have

$$*\omega_2^5\omega_3^5\omega_2^6\omega_3^6\omega_2^7\omega_3^7\omega_2^8\omega_3^8|_{f(S^7)}=\omega_2^1\omega_3^1\omega_4^1\omega_4^5\omega_4^6\omega_4^7\omega_4^8|_{f(S^7)}=\omega_2^1\omega_3^1\omega_4^1\omega_5^1\omega_6^1\omega_7^1\omega_8^1=\mathrm{d}V_{S^7},$$

$$*\omega_2^1\omega_3^1\omega_2^6\omega_3^6\omega_2^7\omega_3^7\omega_2^8\omega_3^8|_{f(S^7)}=-\omega_2^5\omega_3^5\omega_4^5\omega_4^1\omega_4^6\omega_4^7\omega_4^8|_{f(S^7)}=0,$$
$$*\omega_2^5\omega_3^5\omega_2^6\omega_3^6\omega_2^7\omega_4^7\omega_2^8\omega_4^8|_{f(S^7)}=\omega_2^1\omega_3^1\omega_4^1\omega_4^5\omega_4^6\omega_3^7\omega_3^8|_{f(S^7)}=\mathrm{d}V_{S^7},$$
$$*\omega_2^1\omega_3^1\omega_2^6\omega_3^6\omega_2^7\omega_4^7\omega_2^8\omega_4^8|_{f(S^7)}=-\omega_2^5\omega_3^5\omega_4^5\omega_4^1\omega_4^6\omega_3^7\omega_3^8|_{f(S^7)}=0,\ \cdots.$$

Then

$$\frac{1}{b}f^{*}*p_1^2(E)=\frac{2}{5}\cdot\frac{1}{2\pi^4}(3\cdot 3+3\cdot 2)\mathrm{d}V_{S^7}=\frac{3}{\pi^4}\mathrm{d}V_{S^7},$$
$$\frac{2}{5}\int_{S^7}\ f^{*}*p_1^2(E)=1.\quad\square$$

Obviously, we also have $\int_{f(S^7)}\ \tau^{*}\frac{3}{\pi^4}\mathrm{d}V_{S^7}=1$, then

$$[\tau^{*}\frac{3}{\pi^4}\mathrm{d}V_{S^7}]=\frac{2}{5}*p_1^2(E)\in H^7(G(3,8)).$$

Theorem 8.4 (1) The Poincaré dual of $p_1(E)$ is $[M]$ and the Poincaré dual of $\frac{3}{\pi^4}*p_1(E)$ is $[\mathbf{C}P^2]$;

(2) The Poincaré dual of $p_2(F)=p_1^2(E)$ is $[f(S^7)]$ and the Poincaré dual of $\frac{2}{5}*p_1^2(E)$ is $[ASSOC]$.

§9. The case of $G(4,8)$

Let $E=E(4,8),\ F=F(4,8)$ be canonical vector bundles on Grassmann manifold $G(4,8)$. We have

$$p_t(G(4,8))=1+3t^4+4t^8+3t^{12}+t^{16},$$

$$e(E)e(F)=0,\ p_1(E)=-p_1(F),\ p_2(E)=e^2(E),\ p_2(F)=e^2(F),$$

$$p_1^2(E) = p_2(E) + p_2(F),\ p_1(E)p_2(E) = p_1(E)p_2(F) = \frac{1}{2}p_1^3(E),$$
$$p_1^2(E)e(E) = e^3(E),\ p_1^2(F)e(F) = e^3(F).$$

By the method used in §4, we can show $e(E \otimes F) = 6e^4(E) = 6e^4(F)$. Then by $\int_{G(4,8)} e(E \otimes F) = \chi(G(4,8)) = 12$, we have

$$\int_{G(4,8)} e^4(E) = \int_{G(4,8)} e^4(F) = 2.$$

We first study 4 and 12 dimensional cases. As we know $\mathbf{C}P^2, \overline{\mathbf{C}CP}^2, G(2,4)$ can be imbedded in $G(4,8)$ as submanifolds. Under the map $*$: $G(4,8) \to G(4,8)$, $*\mathbf{C}P^2$ is a submanifold of $G(4,8)$. The following table computes the integration of the characteristic classes on the submanifolds of $G(4,8)$.

	$\mathbf{C}P^2$	$*\mathbf{C}P^2$	$G(2,4)$
$e(E)$	0	1	0
$e(F)$	1	0	0
$p_1(E)$	1	-1	2

Note that $\det\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & -1 & 2 \end{pmatrix} = -2$, as proof of Theorem 5.5, we can show $e(E), e(F), p_1(E) \in H^4(G(4,8), \mathbf{Z})$ or $\mathbf{C}P^2, *\mathbf{C}P^2, G(2,4) \in H_4(G(4,8), \mathbf{Z})$ are generators.

By Proposition 2.2, $V(G(4,8)) = \frac{8}{135}\pi^8$ and we can compute

$$a = (e(E), e(E)) = (e(F), e(F)) = \frac{1}{2}(p_1(E), p_1(E)) = \frac{4}{15}\pi^4,$$
$$(e(E), e(F)) = (e(E), p_1(E)) = (e(F), p_1(E)) = 0.$$

In last section we have Cayley submanifold $CAY = \{\pi \in G(4,8) \mid \pi A = -A\}$ of $G(4,8)$. The Lie group $Spin_7$ acts on CAY transitively. Let $e_1, e_2, \cdots, e_8$ be $Spin_7$ frame fields on $\mathbf{R}^8$, then CAY be generated by $e_1e_2e_3e_4$. By the equations listed in the proof of Lemma 7.3, we have

$$\begin{aligned} \mathrm{d}(e_1e_2e_3e_4) &= \sum \omega_i^\alpha E_{i\alpha} \\ &= \omega_1^5(E_{15} + E_{48}) + \omega_1^6(E_{16} + E_{47}) + \omega_1^7(E_{17} - E_{46}) + \omega_1^8(E_{18} - E_{45}) \\ &\quad + \omega_2^5(E_{25} + E_{47}) + \omega_2^6(E_{26} - E_{48}) + \omega_2^7(E_{27} - E_{45}) + \omega_2^8(E_{28} + E_{46}) \\ &\quad + \omega_3^5(E_{35} + E_{46}) + \omega_3^6(E_{36} - E_{45}) + \omega_3^7(E_{37} + E_{48}) + \omega_3^8(E_{38} - E_{47}). \end{aligned}$$

Then the induced metric is

$$\begin{aligned} \mathrm{d}s^2_{CAY} \;=\; & (\omega_1^5-\omega_2^6)^2+(\omega_1^5+\omega_3^7)^2+(\omega_2^6-\omega_3^7)^2 \\ & +(\omega_1^6+\omega_2^5)^2+(\omega_1^6-\omega_3^8)^2+(\omega_2^5-\omega_3^8)^2 \\ & +(\omega_1^7-\omega_3^5)^2+(\omega_1^7-\omega_2^8)^2+(\omega_2^8+\omega_3^5)^2 \\ & +(\omega_1^8+\omega_2^7)^2+(\omega_1^8+\omega_3^6)^2+(\omega_2^7+\omega_3^6)^2. \end{aligned}$$

By $(\omega_1^5-\omega_2^6)(\omega_1^5+\omega_3^7)(\omega_2^6-\omega_3^7)=2\omega_1^5\omega_3^7\omega_2^6,\ \cdots$, we get

$$\mathrm{d}V_{CAY}=16\omega_1^5\omega_2^5\omega_3^5\ \cdots\ \omega_1^8\omega_2^8\omega_3^8.$$

As shown in [7], $2\pi^2 p_1(E)$ is a calibration on $G(4,8)$ with comass $\frac{3}{2}$ and CAY is a calibrated submanifold of $2\pi^2 * p_1(E)$, then

$$2\pi^2 * p_1(E)|_{CAY}=\frac{3}{2}\mathrm{d}V_{CAY}.$$

By triality transformation, we can show that CAY is isometric to $G(3,7)$, then $V(CAY)=V(G(3,7))=\frac{16\pi^6}{45}$. These shows

$$\int_{CAY}\frac{1}{2a}*p_1(E)=\int_{CAY}\frac{45}{32\pi^6}\mathrm{d}V_{CAY}=\frac{1}{2}.$$

By Theorem 3.1, $e(E),e(F),p_1(E)$ can not be generators of $H^4(G(4,8),\mathbf{Z})$. We have proved

Lemma 9.1 $\mathbf{C}P^2, *\mathbf{C}P^2, G(2,4)$ are the generators of $H_4(G(4,8),\mathbf{Z})$, the dual generators are the first column in the following table.

	$\mathbf{C}P^2$	$*\mathbf{C}P^2$	$G(2,4)$
$e(E)$	0	1	0
$e(F)$	1	0	0
$\frac{1}{2}(p_1(E)+e(E)-e(F))$	0	0	1

The inner product of $e(E),e(F),\frac{1}{2}(p_1(E)+e(E)-e(F))$ form a matrix

$$A=\frac{4\pi^4}{15}\begin{pmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & 1 \end{pmatrix},\quad A^{-1}=\frac{15}{4\pi^4}\begin{pmatrix} \frac{3}{2} & -\frac{1}{2} & -1 \\ -\frac{1}{2} & \frac{3}{2} & 1 \\ -1 & 1 & 2 \end{pmatrix}.$$

Lemma 9.2 (1) $*e(E)=\frac{2\pi^4}{15}e^3(E),\ *e(F)=\frac{2\pi^4}{15}e^3(F),\ *p_1(E)=\frac{2\pi^4}{15}p_1^3(E)$;

(2) $\frac{1}{2}(e^3(E)-\frac{1}{2}p_1^3(E)), \frac{1}{2}(e^3(F)+\frac{1}{2}p_1^3(E)), \frac{1}{2}p_1^3(E)\in H^{12}(G(4,8),\mathbf{Z})$ are the generators.

Proof $*e(E), *e(F), *p_1(E)$ and $e^3(E), e^3(F), p_1^3(E)$ are two generators of the coholomogy group $H^{12}(G(4,8))$, they are all the harmonic forms on $G(4,8)$. Then $e^3(E), e^3(F), p_1^3(E)$ can be represented by $*e(E), *e(F), *p_1(E)$. Assuming $e^3(E) = \lambda * e(E) + \mu * e(F) + \nu * p_1(E)$, by $e(F) \wedge *e(E) = 0,\ e(F) \wedge *p_1(E) = 0$, we have $\mu = \nu = 0$,

$$2 = \int_{G(4,8)} e(E) \wedge e^3(E) = \lambda \int_{G(4,8)} e(E) \wedge *e(F) = \lambda \frac{4\pi^4}{15}.$$

Then $*e(E) = \frac{2\pi^4}{15} e^3(E)$. The other two equalities can be proved similarly.

Then

$$\begin{aligned} &(*e(E), *e(F), *\frac{1}{2}(p_1(E) + e(E) - e(F)))A^{-1} \\ &= (\frac{1}{2}(e^3(E) - \frac{1}{2}p_1^3(E)), \frac{1}{2}(e^3(F) + \frac{1}{2}p_1^3(E)), \frac{1}{2}p_1^3(E)). \quad \square \end{aligned}$$

Lemma 9.3 The following table compute the integration of the characteristic classes on the submanifolds of $G(4,8)$.

	$G(4,7)$	$G(3,7)$	CAY
$e^3(E)$	2	0	-1
$e^3(F)$	0	2	1
$p_1^3(E)$	0	0	2

Proof The first column follows from $\int_{G(4,7)} e^3(E(4,8)) = \int_{G(4,7)} e^3(E(4,7)) = 2$ and $e(F(4,8))|_{G(4,7)} = 0,\ p_1^3(E) = 2p_1(E)p_2(F) = 2p_1(E)e^2(F)$. The second column can be proved similarly. For third column, we have proved $\int_{CAY} \frac{1}{2a} *p_1(E) = \frac{1}{2}$, then $\int_{CAY} p_1^3(E) = 2$. By compute $*e(E)|_{CAY},\ *e(F)|_{CAY}$, we can show $\int_{CAY} e^3(E) = -1$ and $\int_{CAY} e^3(F) = 1$. $\square$

Theorem 9.4 (1) $e(E), e(F), \frac{1}{2}(p_1(E) + e(E) - e(F)) \in H^4(G(4,8), \mathbf{Z})$ are the generators, their dual generators are $[\mathbf{C}P^2], [*\mathbf{C}P^2], [G(2,4)] \in H_4(G(4,8), \mathbf{Z})$;

(2) $\frac{1}{2}e^3(E), \frac{1}{2}e^3(F), \frac{1}{2}p_1^3(E)$ and $[G(4,7)], [G(3,7)], [CAY]$ are the generators of $H^{12}(G(4,8), \mathbf{Z})$ and $H_{12}(G(4,8), \mathbf{Z})$ respectively;

(3) The Poincaré duals of $e(E), e(F), \frac{1}{2}(p_1(E) + e(E) - e(F))$ are

$$[G(4,7)],\ [G(3,7)],\ [CAY] + [G(4,7)] - [G(3,7)]$$

respectively.

Proof By Lemma 9.2, $\frac{1}{2}(e^3(E) - \frac{1}{2}p_1^3(E)), \frac{1}{2}(e^3(F) + \frac{1}{2}p_1^3(E)), \frac{1}{2}p_1^3(E)$ are the generators of $H^{12}(G(4,8), \mathbf{Z})$. Then $\frac{1}{2}e^3(E), \frac{1}{2}e^3(F), \frac{1}{2}p_1^3(E)$ are also the generators of $H^{12}(G(4,8), \mathbf{Z})$. $\square$

By Theorem 3.1, we can compute the Poincaré duals of

$$\frac{1}{2}(e^3(E)-\frac{1}{2}p_1^3(E)), \frac{1}{2}(e^3(F)+\frac{1}{2}p_1^3(E)), \frac{1}{2}p_1^3(E).$$

By Theorem 9.4, $\frac{1}{2}(p_1(E)+e(E)-e(F))e(E)=\frac{1}{2}(p_1(E)e(E)+e^2(E))$ and $\frac{1}{2}(p_1(E)e(F)-e^2(F))$, $\frac{1}{2}(p_1(E)e(E)-e^2(E))$, $\frac{1}{2}(p_1(E)e(F)+e^2(F))$ are integral cocycles. The submanifolds $ASSOC$, $A\widetilde{SSOC}$ defined in §7 are also the submanifolds of $G(4,8)$, then $*ASSOC$, $*A\widetilde{SSOC}$ are submanifolds of $G(4,8)$. The following table can be proved by Lemma 7.4.

	$ASSOC$	$A\widetilde{SSOC}$	$*ASSOC$	$*A\widetilde{SSOC}$
$\frac{1}{2}(e^2(F)+p_1(E)e(F))$	1	0	0	0
$\frac{1}{2}(e^2(F)-p_1(E)e(F))$	0	1	0	0
$\frac{1}{2}(e^2(E)+p_1(E)e(E))$	0	0	1	0
$\frac{1}{2}(e^2(E)-p_1(E)e(E))$	0	0	0	1

Theorem 9.5 The characteristic classes

$$\frac{1}{2}(e^2(F)+p_1(E)e(F)),\ \frac{1}{2}(e^2(F)-p_1(E)e(F)),$$

$$\frac{1}{2}(e^2(E)+p_1(E)e(E)),\ \frac{1}{2}(e^2(E)-p_1(E)e(E))$$

are the generators of $H^8(G(4,8),\mathbf{Z})$. Their Poincaré duals are

$$[ASSOC],\ [A\widetilde{SSOC}],\ [*ASSOC],\ [*A\widetilde{SSOC}]$$

respectively.

Proof To see the Poincaré dual of $\xi=\frac{1}{2}(e^2(F)+p_1(E)e(F))$ is $ASSOC$, we want to show that for any $\eta\in H^8(G(4,8))$ we have $\int_{G(4,8)}\xi\wedge\eta=\int_{ASSOC}\eta$. We can take $\eta=\frac{1}{2}(e^2(F)\pm p_1(E)e(F))$, $\frac{1}{2}(e^2(E)\pm p_1(E)e(E))$ to verify this equation. □

By $\mathbf{R}^8=\mathbf{R}^3\oplus\mathbf{R}^5$, we see the product Grassmann $G(2,3)\times G(2,5), G(1,3)\times G(3,5)$ can imbedded in $G(4,8)$ and we have

	$G(4,6)$	$G(2,6)$	$G(2,3)\times G(2,5)$	$G(1,3)\times G(3,5)$
$e^2(E)$	2	0	0	0
$e^2(F)$	0	2	0	0
$p_1(E)e(E)$	0	0	4	0
$p_1(E)e(F)$	0	0	0	4

Then

$$G(4,6),\ G(2,6),\ G(2,3)\times G(2,5),\ G(1,3)\times G(3,5)\in H_8(G(4,8),\mathbf{R})$$

and

$$e^2(E),\ e^2(F),\ p_1(E)e(E),\ p_1(E)e(F)\in H^8(G(4,8),\mathbf{R})$$

are also the generators.

As application, we consider the immersion $f\colon M\to\mathbf{R}^8$ of a compact oriented 4 dimensional manifold, $g\colon M\to G(4,8)$ be its Gauss map. From following table

	$G(4,5)$	$G(1,5)$	$G(2,4)$
$e(E)$	2	0	0
$e(F)$	0	2	0
$p_1(E)$	0	0	2

we have

$$g_*[M]=\frac{1}{2}\chi(M)[G(4,5)]+\lambda[G(1,5)]+\frac{3}{2}Sign(M)[G(2,4)],$$

where $\lambda=\frac{1}{2}\int_M\ e(F(4,8))$ and $Sign(M)$ be the Signature of M. $\lambda=0$ if f is an imbedding.

If g is the Gauss map of immersion of in $\mathbf{R}^7$ or $\mathbf{R}^6$, we have

$$g_*[M]=\frac{1}{2}\chi(M)[G(4,5)]+\frac{3}{2}Sign(M)[G(2,4)].$$

§10. The cohomology groups on *ASSOC*

The submanifold $ASSOC\approx G_2/SO(4)$ of Grassmann manifold $G(3,7)$ is important in theory of calibration, see [7], [9]. In [4] Borel and Hirzebruch studied the characteristic classes on homogenous spaces, they computed the cohomology of $ASSOC$. In the following we use Gysin sequence to study the cohomology of $ASSOC$.

As §7, let $G(2,7)$ and $G(3,7)$ be Grassmann manifolds on $\mathbf{R}^7\subset\mathbf{R}^8$ generated by $\bar{e}_2,\cdots,\bar{e}_8$, and $S^6\subset\mathbf{R}^7$ the unit sphere. There is a fibre bundle $\tau_1\colon G(2,7)\to S^6$ defined by $\pi A=\bar{e}_1vA$, $\tau_1(\pi)=v$, where $A\in C\ell_8$ is defined in §7. For any $G\in G_2$, we have the following commutative diagram

$$\begin{tikzcd} G(2,7) \arrow[r, "G"] \arrow[d, "\tau_1"'] & G(2,7) \arrow[d, "\tau_1"] \\ S^6 \arrow[r, "G"] & S^6. \end{tikzcd}$$

The fibre $\tau_1^{-1}(\bar{e}_2) = \{v \wedge Jv \mid v \in S^6,\ v \perp \bar{e}_2\} \approx \mathbf{C}P^2$, see [16].

Then for any $\pi \in G(2,7),\ v = \tau_1(\pi),\ v \wedge \pi \in ASSOC$. This defines map

$$\tau_2\colon\ G(2,7) \to ASSOC, \quad \pi \mapsto v \wedge \pi.$$

For any $e_2e_3e_4 \in ASSOC,\ e_2e_3e_4A = \bar{e}_1A$, then $\tau_2(e_3e_4) = e_2e_3e_4$. This shows

Lemma 10.1 $\tau_2\colon\ G(2,7) \to ASSOC$ is a fibre bundle with fibre $G(2,3) = S^2$.

Let $i\colon\ ASSOC \to G(3,7)$ be inclusion. It is easy to see $G(2,7)$ isomorphic to the sphere bundle $S(\tilde{E}) = \{v \in \tilde{E},\ |v| = 1\}$ of the induced bundle $\tilde{E} = i^*E(3,7)$. Let $e(E(3,7)) \in H^3(G(3,7),\mathbf{Z})$ be the Euler class of $E(3,7)$, $2e(E(3,7)) = 0$, see [12] p.95-103. Then $e(\tilde{E}) = i^*e(E(3,7)) \in H^3(ASSOC,\mathbf{Z})$ is the Euler class of the induced bundle $\tilde{E}$. There is a Gysin exact sequence for the sphere bundle $G(2,7) \to ASSOC$,

$$\begin{aligned} &\longrightarrow H^q(ASSOC) \xrightarrow{\tau_2^*} H^q(G(2,7)) \xrightarrow{\tau_{2*}} H^{q-2}(ASSOC) \\ &\xrightarrow{\wedge e(\tilde{E})} H^{q+1}(ASSOC) \xrightarrow{\tau_2^*} H^{q+1}(G(2,7)) \longrightarrow, \end{aligned}$$

where τ_{2*} is the integration along the fibre. The coefficients of the cohomology groups can be $\mathbf{R}, \mathbf{Z}$ or $\mathbf{Z}_2$.

Lemma 10.2 $e(\tilde{E}) = i^*e(E(3,7)) \neq 0$.

Proof The map $\tau_{2*}\colon\ H^q(G(2,7),\mathbf{Z}) \to H^{q-2}(ASSOC,\mathbf{Z})$ is the integration along the fibre. Let $\bar{e}_1, e_2, e_3, \cdots, e_8$ be G_2 frame fields, $G(2,7)$ is generated by e_3e_4 and $\tau_2(e_3e_4) = e_2e_3e_4$. Then the Euler class of vector bundle $E(2,7)$ can be represented by

$$e(E(2,7)) = \frac{1}{2\pi}\omega_3^2 \wedge \omega_4^2 + \frac{1}{2\pi}\sum_{\alpha=5}^{8} \omega_3^\alpha \wedge \omega_4^\alpha$$

and $\omega_3^2 \wedge \omega_4^2$ is the volume element of the fibre at e_3e_4. Then $\tau_{2*}(e(E(2,7)) = 2$.

By Gysin sequence, the map $\tau_{2*}\colon\ H^2(G(2,7)) \to H^0(ASSOC)$ is surjective if $e(\tilde{E}) = 0$. These contradicts to the fact $\tau_{2*}(e(E(2,7)) = 2$ and $e(E(2,7)) \in H^2(G(2,7),\mathbf{Z})$ be a generator. $\Box$

Then $e(E(3,7))$ is also nonzero and a torsion element of $H^3(G(3,7),\mathbf{Z})$.

Theorem 10.3 The cohomology groups of $ASSOC$ are

$$H^q(ASSOC,\mathbf{Z}_2) = \begin{cases} \mathbf{Z}_2, & q \neq 1,7, \\ 0, & q = 1,7; \end{cases}$$

$$H^q(ASSOC,\mathbf{Z}) = \begin{cases} \mathbf{Z}, & q = 0,4,8, \\ \mathbf{Z}_2, & q = 3,6, \\ 0, & q = 1,2,5,7; \end{cases}$$

$$H^q(ASSOC, \mathbf{R}) = \begin{cases} \mathbf{R}, & q = 0, 4, 8, \\ 0, & q \neq 0, 4, 8. \end{cases}$$

Proof $G(2,7)$ is a Kaehler manifold, the coholomogy of $G(2,7)$ is generated by Euler class $e(E(2,7))$. We prove the case of $\mathbf{Z}_2$ coefficients, the other cases are left to the reader.

By Gysin sequence, we have

$$0 = H^{-2}(ASSOC) \xrightarrow{\wedge e(\tilde{E})} H^1(ASSOC) \xrightarrow{\tau_2^*} H^1(G(2,7)) = 0,$$

$$0 = H^{-1}(ASSOC) \xrightarrow{\wedge e(\tilde{E})} H^2(ASSOC) \xrightarrow{\tau_2^*} H^2(G(2,7)) \xrightarrow{\tau_{2*}=0} H^0(ASSOC)$$
$$\xrightarrow{\wedge e(\tilde{E})} H^3(ASSOC) \xrightarrow{\tau_2^*} H^3(G(2,7)) = 0.$$

These shows $H^1(ASSOC) = 0$ and $H^2(ASSOC) \cong H^2(G(2,7))$, $H^0(ASSOC) \cong H^3(ASSOC)$. By

$$0 = H^1(ASSOC) \xrightarrow{\wedge e(\tilde{E})} H^4(ASSOC) \xrightarrow{\tau_2^*} H^4(G(2,7)) \xrightarrow{\tau_{2*}} H^2(ASSOC)$$

and $\tau_{2*} = 0$: $H^4(G(2,7), \mathbf{Z}_2) \to H^4(ASSOC, \mathbf{Z}_2)$, we have

$$H^4(ASSOC) \cong H^4(G(2,7)).$$

The cases of $q = 5, \cdots, 8$ can be proved similarly or by Poincaré duality.

References

[1] Chen, W.H., The differential geometry of Grassmann manifold as submanifold, Acta Math., Sinica, **31A**(1988), 46-53.

[2] Chern, S.S., Characteristic classes of Hermitian manifolds. Annals of. Math. **47**(1946), 85-121.

[3] Chern, S.S. and Spanier, E.H., A theorem on orientable surfaces in four-dimensional space. Comm. Math. Helv. **25**(1951), 205-209.

[4] Borel, A. and Hirzebruch, F., Characteristic classes and homogenous spaces, I. Amer. J. Math. **80**(1958), 458-538.

[5] Bott, R. and Tu, L.W., Differential forms in algebraic topology, Springer-Verlag, New York, GTM **82**, 1982.

[6] Bryant, R.L., Submanifolds and special structures on the octonians, Diff Geom. J., **17**(1982), 185-232.

[7] Gluck, H., Mackenzie, D. and Morgan, F., Volume-minimizing cycles in Grassmann manifolds, Duke Math. J., **79**(1995), 335-404.

[8] Greub, W., Halperin, S. and Vanstone, R., Connections, curvature, and cohomology, Academic Press, New Yark, 1976.

[9] Harvey, F.R., and Lawson, Jr H.B., Calibrated Geometries, Acta Math. J., **148**(1982), 47-157.

[10] Lawson, Jr H.B., Michelsohn, M., Spin geometry. Princeton University Press. Princeton New Jersey, 1989

[11] Kabayashi, S. and Nomizu, K., Foundations of differential geometry, vol. 2, Interscience Publishers, New York, 1969.

[12] Milnor, J.W. and Stasheff, J.D., Characteristic classes, Ann. Math. Studies., No. **76**, Princeton, 1974.

[13] Wolf, J.A., Spaces of constant curvature, McGraw-Hill, New Yark, 1967.

[14] Zhou, J.W., Irreducible Clifford Modules, Tsukuba J. Math., **27**(2003), 57-75.

[15] Zhou, J.W., Spinors, Calibrations and Grassmannians, Tsukuba J. Math., **27**(2003), 77-97.

[16] Zhou, J.W. and Huang, H., Geometry on Grassmann Manifolds G(2,8) and G(3,8), Math. J. of Okayama Univ., **44**(2002), 171-179.

[17] Zhou, J.W., Morse functions on Grassmann manifolds, Proc. of the Royal Soc. of Edinburgh, **135A**(2005), 1-13.

[18] Zhou, J.W., Totally geodesic submanifolds in Lie groups, Acta Math. Hungarica. **111**(2006), 29-41.

[19] Zhou, J.W., The geometry and topology on Grassmann manifolds, Math. J. Okayama Univ,. **48**(2006), 181-195.

[20] Zhou, J.W., A note on characteristic classes, Czechoslovak Math. J., 562006721-732.